arXiv:1107.3654v2 [math.AG] 2 Dec 2011

STANDARD MONOMIAL THEORY FOR DESINGULARIZED RICHARDSON VARIETIES IN THE FLAG VARIETY $GL(n)/B$

MICHAËL BALAN

Abstract. We consider a desingularization Γ of a Richardson variety $X_w^v = X_w \cap X^v$ in the flag variety $F\ell(n) = GL(n)/B$, obtained as a fibre of a projection from a certain Bott-Samelson variety Z. We then construct a basis of the homogeneous coordinate ring of Γ inside Z, indexed by combinatorial objects which we call w_0-*standard tableaux*.

Introduction

Standard Monomial Theory (SMT) originated in the work of Hodge [19], who considered it in the case of the Grassmannian $G_{d,n}$ of d-subspaces of a (complex) vector space of dimension n. The homogeneous coordinate ring $\mathbf{C}[G_{d,n}]$ is the quotient of the polynomial ring in the Plücker coordinates $p_{i_1 \ldots i_d}$ by the Plücker relations, and Hodge provided a combinatorial rule to select, among all monomials in the $p_{i_1 \ldots i_d}$, a subset that forms a basis of $\mathbf{C}[G_{d,n}]$: these (so-called standard) monomials are parametrized by semi-standard Young tableaux. Moreover, he showed that this basis is compatible with any Schubert variety $X \subset G_{d,n}$, in the sense that those basis elements that remain non-zero when restricted to X can be characterized combinatorially, and still form a basis of $\mathbf{C}[X]$. The aim of SMT is then to generalize Hodge's result to any flag variety G/P (G a connected semi-simple group, P a parabolic subgroup): in a more modern formulation, the problem consists, given a line bundle L on G/P, in producing a "nice" basis of the space of sections $H^0(X, L)$ ($X \subset G/P$ a Schubert variety), parametrized by some combinatorial objects. SMT was developed by Lakshmibai and Seshadri (see [29, 30]) for groups of classical type, and Littelmann extended it to groups of arbitrary type (including in the Kac-Moody setting), using techniques such as the path model in representation theory [32, 33] and Lusztig's Frobenius map for quantum groups at roots of unity [34]. Standard Monomial Theory has numerous applications in the geometry of Schubert varieties: normality, vanishing theorems, ideal theory, singularities, and so on [26].

Richardson varieties, named after [36], are intersections of a Schubert variety and an opposite Schubert variety inside a flag variety G/P. They previously appeared in [20, Ch. XIV, §4] and [38], as well as the corresponding open subvarieties in [10]. They have since played a role in different contexts, such as equivariant K-theory [25], positivity in Grothendieck groups [5], standard monomial theory [7], Poisson geometry [13], positroid varieties [21], and their generalizations [22, 2]. In particular, SMT on G/P is known to be compatible with Richardson varieties [25] (at least for a very ample line bundle on G/P).

Date: December 2, 2011.

2010 *Mathematics Subject Classification.* Primary 14M15, Secondary 05E10 14M17 20G05.

Like Schubert varieties, Richardson varieties may be singular [24, 23, 40, 1]. Desingularizations of Schubert varieties are well known: they are the Bott-Samelson varieties [4, 9, 14], which are also used for example to establish some properties of Schubert polynomials [35], or to give criteria for the smoothness of Schubert varieties [12, 8]. An SMT has been developed for Bott-Samelson varieties of type A_n in [28], and of arbitrary type in [27] using the path model [32, 33].

In the present paper, we shall describe a Standard Monomial Theory for a desingularization of a Richardson variety. To be more precise, we introduce some notations. Let $G = GL(n, k)$ where k is an algebraically closed field of arbitrary characteristic, B the Borel subgroup of upper triangular matrices, and $T \subset B$ the maximal torus of diagonal matrices. The quotient G/B identifies with the variety $F\ell(n)$ of all complete flags in k^n. Let $(e_1, \dots, e_n)$ be the canonical basis of k^n. To each permutation $w \in S_n$, we can associate a T-fixed point e_w in $F\ell(n)$: its ith constituent is the space generated by $e_{w(1)}, \dots, e_{w(i)}$. We denote by F_{can} the T-fixed point corresponding to the identity e of S_n, and $F_{\mathrm{op\,can}}$ the T-fixed point e_{w_0}, where w_0 is the longest element of S_n. The symmetric group S_n is generated by the simple transpositions $s_i = (i, i+1)$, $i = 1, \dots, n$. We denote a permutation $u \in S_n$ with the one-line notation $[u(1)\ u(2) \dots\ u(n)]$. Denote by B^- the subgroup of G of lower triangular matrices and consider the Schubert cells $C_w = B.e_w$ and the opposite Schubert cells $C^v = B^-.e_v$. The Richardson variety $X_w^v \subset F\ell(n)$ is the intersection of the direct Schubert variety $X_w = \overline{C_w}$ with the opposite Schubert variety $X^v = \overline{C^v} = w_0 X_{w_0 v}$. Fix a reduced decomposition $w = s_{i_1} \dots s_{i_d}$ and consider the Bott-Samelson desingularization $Z = Z_{i_1 \dots i_d}(F_{\mathrm{can}}) \to X_w$, and similarly $Z' = Z_{i_r i_{r-1} \dots i_{d+1}}(F_{\mathrm{op\,can}}) \to X^v$ for a reduced decomposition $w_0 v = s_{i_r} s_{i_{r-1}} \dots s_{i_{d+1}}$. Then the fibred product $Z \times_{F\ell(n)} Z'$ has been considered as a desingularization of X_w^v in [6], but for our purposes, it will be more convenient to realize it as the fibre $\Gamma_{\mathbf{i}}$ ($\mathbf{i} = i_1 \dots i_d i_{d+1} \dots i_r$) of the projection $Z_{\mathbf{i}} = Z_{\mathbf{i}}(F_{\mathrm{can}}) \to F\ell(n)$ over $F_{\mathrm{op\,can}}$ (see Section 1 for the precise connection between those two constructions).

In [28, 27], Lakshmibai, Littelmann, and Magyar define a family of line bundles $L_{\mathbf{i},\mathbf{m}}$ ($\mathbf{m} = m_1 \dots m_r \in \mathbf{Z}^r_{\geq 0}$) on $Z_{\mathbf{i}}$ (they are the only globally generated line bundles on $Z_{\mathbf{i}}$, as pointed out in [31]), and give a basis for the space of sections $H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$. In [28], the elements p_T of this basis, called standard monomials, are indexed by combinatorial objects T called standard tableaux: the latter's definition involves certain sequences $J_{11} \supset \cdots \supset J_{1m_1} \supset \cdots \supset J_{r1} \supset \cdots \supset J_{rm_r}$ of subwords of $\mathbf{i}$, called liftings of T (see Section 2 for precise definitions—actually, two equivalent definitions of standard tableaux are given in [28], but we will only use the one in terms of liftings). Note also that $L_{\mathbf{i},\mathbf{m}}$ is very ample precisely when $m_j > 0$ for all j (see [31], Theorem 3.1), in which case $\mathbf{m}$ is called regular.

The main result of this paper states that in this case, SMT on $Z_{\mathbf{i}}$ is compatible with $\Gamma_{\mathbf{i}}$.

Theorem 0.1. *Assume that* $\mathbf{m}$ *is regular. With the above notation, the standard monomials* p_T *such that* $(p_T)_{|\Gamma_{\mathbf{i}}} \neq 0$ *still form a basis of* $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$.
Moreover, $(p_T)_{|\Gamma_{\mathbf{i}}} \neq 0$ *if and only if* T *admits a lifting* $J_{11} \supset \cdots \supset J_{rm_r}$ *such that each subword* J_{km} *contains a reduced expression of* w_0.

We prove this theorem in three steps.

1. Call T (or p_T) w_0-standard if the above condition on (J_{km}) holds. We prove by induction over $M = \sum_{j=1}^{r} m_j$ that the w_0-standard monomials p_T are linearly independent on $\Gamma_{\mathbf{i}}$. (Here the assumption that $\mathbf{m}$ is regular is not necessary.)
2. In the regular case, we prove that a standard monomial p_T does not vanish identically on $\Gamma_{\mathbf{i}}$ if and only if it is w_0-standard, using the combinatorics of the Demazure product (see Definition 4.2). It follows that w_0-standard monomials form a basis of the homogeneous coordinate ring of $\Gamma_{\mathbf{i}}$ (when $\Gamma_{\mathbf{i}}$ is embedded in a projective space via the very ample line bundle $L_{\mathbf{i},\mathbf{m}}$).
3. We use cohomological techniques to prove that the restriction map
$$H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$$
is surjective. More explicitly, we define a family $(Y_{\mathbf{i}}^{u})$ of subvarieties of $Z_{\mathbf{i}}$ indexed by S_n, with the property that $Y_{\mathbf{i}}^{e} = Z_{\mathbf{i}}$ and $Y_{\mathbf{i}}^{w_0} = \Gamma_{\mathbf{i}}$. We construct a sequence in S_n, $e = u_0 < u_1 < \cdots < u_N = w_0$, such that for every t, $Y_{\mathbf{i}}^{u_{t+1}}$ is defined in $Y_{\mathbf{i}}^{u_t}$ by the vanishing of a single Plücker coordinate p_κ, in such a way that each restriction map $H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}})$ can be shown to be surjective using vanishing theorems (Corollary 5.7 and Theorem 5.23). This shows that the w_0-standard monomials span $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$.

Note that alternate bases for certain Bott-Samelson varieties have been constructed in [39], and the fibred products $Z \times_{F\ell(n)} Z'$ have been studied from this point of view in [11].

Sections are organized as follows: in Section 1, we first fix notation and recall information on Bott-Samelson varieties $Z_{\mathbf{i}}$, and then show that the fibre $\Gamma_{\mathbf{i}}$ of $Z_{\mathbf{i}} \to F\ell(n)$ over $F_{\mathrm{op\,can}}$ is a desingularization of the Richardson variety X_w^v; this fact is most certainly known to experts, but has not, to our knowledge, appeared in the literature. In Section 2, we recall the main results about SMT for Bott-Samelson varieties from [28], in particular the definition of standard tableaux. In Section 3, we define w_0-standard monomials and we prove that they are linearly independent in $\Gamma_{\mathbf{i}}$. In Section 4, we prove that when $\mathbf{m}$ is regular, a standard monomial does not vanish identically on $\Gamma_{\mathbf{i}}$ if and only if it is w_0-standard. Finally, we prove in Section 5 that w_0-standard monomials generate the space of sections $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$.

Acknowledgements. I would like to thank Christian Ohn for helpful discussions. I am also grateful to Michel Brion for his valuable remarks, and especially for pointing out gaps in the proofs of Proposition 5.16 and Theorem 5.23.

1. Desingularized Richardson varieties

The notations are as in the Introduction. In addition, if $k, l \in \mathbf{Z}$, then we denote by $[k, l]$ the set $\{k, k+1, \dots, l\}$, and by $[l]$ the set $[1, l]$.

We first recall a number of results on Bott-Samelson varieties (see *e.g.* [35]).

Definition 1.1. Two flags F, G in $F\ell(n)$ are called *i-adjacent* if they coincide except (possibly) at their components of dimension i, a situation denoted by $F \overset{i}{—} G$.

Notations 1.2. For $i \in [n]$, we denote by $F\ell(\hat{i})$ the variety of partial flags

$$V_1 \subset V_2 \subset \cdots \subset V_{i-1} \subset V_{i+1} \subset \cdots \subset V_n, \ (\dim V_j = j),$$

and by $\psi_{\hat{i}} : F\ell(n) \to F\ell(\hat{i})$ the natural projection.

Then F and G are i-adjacent if and only if they have the same image by $\psi_{\hat{i}}$.

Consider a word $\mathbf{i} = i_1 \dots i_r$ in $[n-1]$, with $w(\mathbf{i}) = s_{i_1} \dots s_{i_r} \in S_n$ not necessarily reduced. A *gallery of type* $\mathbf{i}$ is a sequence of the form

$$F_0 \overset{i_1}{\text{---}} F_1 \overset{i_2}{\text{---}} \dots \overset{i_r}{\text{---}} F_r. \tag{1}$$

For a given flag F_0, the *Bott-Samelson variety* of type $\mathbf{i}$ starting at F_0 is the set of all galleries (1), *i.e.* the fibred product

$$Z_{\mathbf{i}}(F_0) = \{F_0\} \times_{F\ell(\hat{i}_1)} F\ell(n) \times_{F\ell(\hat{i}_2)} \cdots \times_{F\ell(\hat{i}_r)} F\ell(n)$$

(a subvariety of $F\ell(n)^r$). In particular, $Z_{i_1 \dots i_r}(F_0)$ is a $\mathbf{P}^1$-fibration over $Z_{i_1 \dots i_{r-1}}(F_0)$, which shows by induction over r that Bott-Samelson varieties are smooth.

Each subset $J = \{j_1 < \cdots < j_k\} \subset [r]$ defines a subword $\mathbf{i}(J) = (i_{j_1}, \dots, i_{j_k})$ of $\mathbf{i}$. We then write $Z_J(F_0)$ instead of $Z_{\mathbf{i}(J)}(F_0)$, and we view it as the subvariety of $Z_{\mathbf{i}}(F_0)$ consisting of all galleries (1) such that $F_{j-1} = F_j$ whenever $j \notin J$.

We denote by $F_{\mathrm{can}} : \langle e_1 \rangle \subset \langle e_1, e_2 \rangle \subset \cdots \subset \langle e_1, e_2, \dots, e_n \rangle$ the flag associated to the canonical basis, and by $F_{\mathrm{op\,can}} : \langle e_n \rangle \subset \langle e_n, e_{n-1} \rangle \subset \cdots \subset \langle e_n, e_{n-1}, \dots, e_1 \rangle$ the opposite canonical flag. Note that $F_{\mathrm{op\,can}} = e_{w_0}$.

In the sequel, we shall only need galleries starting at F_{can} or at $F_{\mathrm{op\,can}}$; in particular, we write $Z_{\mathbf{i}} = Z_{\mathbf{i}}(F_{\mathrm{can}})$.

The (diagonal) B-action on $F\ell(n)^r$ leaves $Z_{\mathbf{i}}$ invariant. In particular, the T-fixed points of $Z_{\mathbf{i}}$ are the galleries of the form

$$F_{\mathrm{can}} \overset{i_1}{\text{---}} e_{u_1} \overset{i_2}{\text{---}} e_{u_1 u_2} \overset{i_3}{\text{---}} \dots \overset{i_r}{\text{---}} e_{u_1 \dots u_r},$$

where each $u_j \in S_n$ is either e or s_{i_j}. This gallery will be denoted $e_J \in Z_{\mathbf{i}}$, where $J = \{j \mid u_j = s_{i_j}\} = \{j_1 < \cdots < j_k\}$.

For $j \in [r]$, we denote by $\mathrm{pr}_j : Z_{\mathbf{i}} \to F\ell(n)$ the projection sending the gallery (1) to F_j. Note that $w(\mathbf{i}(J)) = s_{i_{j_1}} \dots s_{i_{j_k}} = u_1 \dots u_r$, so $\mathrm{pr}_r(e_J) = e_{u_1 \dots u_r} = e_{w(\mathbf{i}(J))}$.

When $\mathbf{i}$ is reduced, *i.e.* $w = s_{i_1} \dots s_{i_r}$ is a reduced expression in S_n, a flag F lies in the Schubert variety X_w if and only if there is a gallery of type $\mathbf{i} = i_1 \dots i_r$ from F_{can} to F, hence the last projection pr_r takes $Z_{\mathbf{i}}$ surjectively to X_w. Moreover, this surjection is birational: it restricts to an isomorphism over the Schubert cell C_w: thus, $\mathrm{pr}_r : Z_{\mathbf{i}} \to X_w$ is a desingularization of X_w, and likewise for the last projection $Z_{\mathbf{i}}(F_{\mathrm{op\,can}}) \to X^{w_0 w}$.

When $\mathbf{i}$ is not necessarily reduced, $\mathrm{pr}_r(Z_{\mathbf{i}})$ may be described as follows. Recall [28, Definition-Lemma 1] that the poset $\{w(\mathbf{i}(J)) \mid J \subset [r]\}$ admits a unique maximal element, denoted by $w_{\max}(\mathbf{i})$ (so $w_{\max}(\mathbf{i}) = w(\mathbf{i})$ if and only if $\mathbf{i}$ is reduced):

Proposition 1.3. *Let* $\mathbf{i}$ *be an arbitrary word. Then* $\mathrm{pr}_r(Z_{\mathbf{i}})$ *is the Schubert variety* X_w*, where* $w = w_{max}(\mathbf{i})$.

Proof. Since $\mathrm{pr}_r(Z_{\mathbf{i}})$ is B-stable, it is a union of Schubert cells. But $Z_{\mathbf{i}}$ is a projective variety, so the morphism pr_r is closed, hence $\mathrm{pr}_r(Z_{\mathbf{i}})$ is a union of Schubert varieties, and therefore a single Schubert variety X_w since $Z_{\mathbf{i}}$ is irreducible. Moreover, the T-fixed points e_J in $Z_{\mathbf{i}}$ project to the T-fixed points $e_{w(\mathbf{i}(J))}$ in X_w, and all T-fixed points of X_w are obtained in this way (indeed, if e_v is such a point, then the fibre $\mathrm{pr}_r^{-1}(e_v)$ is T-stable, so it must contain some e_J by Borel's fixed point theorem). In particular, e_w corresponds to a choice of $J \subset \{1, \ldots, r\}$ such that $w(\mathbf{i}(J))$ is maximal, hence the result. $\square$

We now turn to the description of a desingularization of a Richardson variety $X_w^v = X_w \cap X^v$, $v \leq w \in S_n$. Let $Z = Z_{i_1 \ldots i_d}$ for some reduced decomposition $w = s_{i_1} \ldots s_{i_d}$ and $Z' = Z_{i_r \ldots i_{d+1}}(F_{\mathrm{op\,can}})$ for some reduced decomposition $w_0 v = s_{i_r} s_{i_{r-1}} \ldots s_{i_{d+1}}$. Since Z desingularizes X_w and Z' desingularizes X^v, a natural candidate for a desingularization of X_w^v is the fibred product $Z \times_{F\ell(n)} Z'$. However, we wish to see this variety in a slightly different way: an element of $Z \times Z'$ is a pair of galleries

$$F_{\mathrm{can}} \overset{i_1}{—} F_1 \overset{i_2}{—} \ldots \overset{i_d}{—} F_d,$$

$$F_{\mathrm{op\,can}} \overset{i_r}{—} G_{r-1} \overset{i_{r-1}}{—} \ldots \overset{i_{d+1}}{—} G_d,$$

and it belongs to $Z \times_{F\ell(n)} Z'$ when the end points F_d and G_d coincide: in this case, by reversing the second gallery, they concatenate to form a longer gallery

$$F_{\mathrm{can}} \overset{i_1}{—} F_1 \overset{i_2}{—} \ldots \overset{i_d}{—} F_d \overset{i_{d+1}}{—} \ldots \overset{i_r}{—} F_{\mathrm{op\,can}}.$$

Thus, $Z \times_{F\ell(n)} Z'$ identifies with the set of all galleries in $Z_{\mathbf{i}} = Z_{i_1 \ldots i_r}$ that end in $F_{\mathrm{op\,can}}$, *i.e.* with the fibre

$$\Gamma_{\mathbf{i}} = \mathrm{pr}_r^{-1}(F_{\mathrm{op\,can}})$$

of the last projection $\mathrm{pr}_r : Z_{\mathbf{i}} \to F\ell(n)$. By construction, the dth projection pr_d then maps $\Gamma_{\mathbf{i}}$ onto the Richardson variety X_w^v.

Proposition 1.4. *In the above notation, the dth projection* $\mathrm{pr}_d : \Gamma_{\mathbf{i}} \to X_w^v$ *is a desingularization,* i.e. pr_d *is birational, and the variety* $\Gamma_{\mathbf{i}}$ *is smooth and irreducible.*

Proof. We first compute the dimension of $\Gamma_{\mathbf{i}}$: since pr_r is surjective, there exists a non-empty open set O in $F\ell(n)$ such that every point $F \in O$ has a fibre of pure dimension $\dim(Z_{\mathbf{i}}) - \dim(F\ell(n))$. Since the flag variety $F\ell(n)$ is irreducible, O meets the open Schubert cell C_{w_0}. Let $F \in O \cap C_{w_0}$. Since pr_r is B-equivariant, the fibres of F and $F_{\mathrm{op\,can}} = e_{w_0}$ are isomorphic. In particular, they have the same dimension, so $\dim(\Gamma_{\mathbf{i}}) = \dim(Z_{\mathbf{i}}) - \dim(F\ell(n))$.

Next we show that $\Gamma_{\mathbf{i}}$ is smooth. Let $\gamma \in \Gamma_{\mathbf{i}}$. We want to prove that the tangent space $T_\gamma(\Gamma_{\mathbf{i}})$ of $\Gamma_{\mathbf{i}}$ at γ and $\Gamma_{\mathbf{i}}$ have the same dimension. Let $\Omega = \mathrm{pr}_r^{-1}(C_{w_0})$. Let U be the maximal unipotent subgroup of B. This subgroup acts simply transitively on the Schubert cell C_{w_0}. Consider the morphism

$$\begin{array}{rrcl} s: & C_{w_0} = U.e_{w_0} & \to & \Omega \\ & u.e_{w_0} & \mapsto & u.\gamma \end{array}$$

Since pr_r is U-equivariant, we have $\mathrm{pr}_r \circ s = \mathrm{id}_{C_{w_0}}$. Differentiating this equality in e_{w_0} gives $\mathrm{d}\,\mathrm{pr}_r(\gamma) \circ \mathrm{d}\,s(e_{w_0}) = \mathrm{id}_{T_{e_{w_0}} F\ell(n)}$. In particular, the linear map $\mathrm{d}\,\mathrm{pr}_r(\gamma) : T_\gamma(Z_{\mathbf{i}}) \to T_{e_{w_0}}(F\ell(n))$ is surjective. Moreover, $T_\gamma(\Gamma_{\mathbf{i}}) \subset \ker(\mathrm{d}\,\mathrm{pr}_r(\gamma))$. From this, we deduce

$$\begin{aligned} \dim(\Gamma_{\mathbf{i}}) \leq \dim T_\gamma(\Gamma_{\mathbf{i}}) &\leq \dim T_\gamma(Z_{\mathbf{i}}) - \dim T_{e_{w_0}}(F\ell(n)) \\ &\leq \dim Z_{\mathbf{i}} - \dim F\ell(n) \text{ (since } Z_{\mathbf{i}} \text{ and } F\ell(n) \text{ are both smooth)} \\ &\leq \dim \Gamma_{\mathbf{i}}, \end{aligned}$$

hence $\Gamma_{\mathbf{i}}$ is smooth.

Now we show that $\Gamma_{\mathbf{i}}$ is irreducible. Let $C_1, \dots, C_e$ be the irreducible components of $\Gamma_{\mathbf{i}}$. Since $\Gamma_{\mathbf{i}}$ is smooth, the C_j are also the connected components of $\Gamma_{\mathbf{i}}$. The variety Ω is open in $Z_{\mathbf{i}}$. In particular, Ω is irreducible. Since pr_r is B-equivariant,

$$\Omega = \bigcup_{i=1}^{e} \bigcup_{b \in B} bC_i.$$

Let $\Omega_i = \bigcup_{b\in B} bC_i$. The morphism $f : U \times \Gamma_{\mathbf{i}} \to \Omega, (b, \gamma) \mapsto b.\gamma$ is an isomorphism. In particular, $\Omega_i = f(U \times C_i)$ is an irreducible closed set in Ω. So $\Omega = \bigcup_{i=1}^{e} \Omega_i$ is a disjoint decomposition of Ω into irreducibles. Hence $e = 1$, and $\Gamma_{\mathbf{i}}$ is irreducible.

Finally, to show that $\Gamma_{\mathbf{i}} \to X_w^v$ is birational, we consider the projections $\mathrm{pr}_d : Z \to X_w$ and $\mathrm{pr}_{r-d} : Z' \to X^v$. Since they are birational, there exist open subsets $U_w \subset X_w$ and $O \subset Z$ isomorphic under pr_d, and open subsets $U^v \subset X^v$ and $O' \subset Z'$ isomorphic under pr_{r-d}. Then the open set $(O \times O') \cap (Z \times_{F\ell(n)} Z')$ of $Z \times_{F\ell(n)} Z'$ is isomorphic to the open set $U_w \cap U^v$ of X_w^v under $\mathrm{pr}_d : Z \times_{F\ell(n)} Z' \to X_w^v$. Since X_w^v and $Z \times_{F\ell(n)} Z' \cong \Gamma_{\mathbf{i}}$ are irreducible, these open subsets must be dense. The birationality of $\mathrm{pr}_d : \Gamma_{\mathbf{i}} \to X_w^v$ follows. □

Remark 1.5. In characteristic 0, it can be proved more directly that the fibred product $Z \times_{F\ell(n)} Z'$ is smooth using Kleiman's transversality theorem (*cf.* [16], Theorem 10.8). Moreover, this theorem also states that every irreducible component of $Z \times_{F\ell(n)} Z'$ is of dimension $\dim(Z) + \dim(Z') - \dim(F\ell(n))$. To prove the irreducibility of $Z \times_{F\ell(n)} Z'$, consider ∂Z (resp. $\partial Z'$) the union of all Bott-Samelson varieties X with $X \subsetneq Z$ (resp. $X \subsetneq Z'$). By Kleiman's transversality theorem, the dimension of $(\partial Z \times_{F\ell(n)} Z') \cup (Z \times_{F\ell(n)} \partial Z')$ is less than $\dim(Z \times_{F\ell(n)} Z')$. So, on one hand, the fibred-product $O = (Z \setminus \partial Z) \times_{F\ell(n)} (Z' \setminus \partial Z')$ meets each irreducible component of $Z \times_{F\ell(n)} Z'$, hence O is dense. On the other hand, O is isomorphic to the open Richardson variety $C_w^v = C_w \cap C^v$, hence O is irreducible. Therefore, $Z \times_{F\ell(n)} Z'$ is irreducible.

For $\mathbf{i}$ an arbitrary word, we may still consider the variety $\Gamma_{\mathbf{i}}$ of galleries of type $\mathbf{i}$, beginning at F_{can} and ending at $F_{\mathrm{op\,can}}$. In general this variety is no longer birational to a Richardson variety. But we still have

Proposition 1.6. *Let $\mathbf{i} = i_1 \dots i_r$ be an arbitrary word, and consider the projection $\mathrm{pr}_j : \Gamma_{\mathbf{i}} \to F\ell(n)$. Then $\mathrm{pr}_j(\Gamma_{\mathbf{i}})$ is the Richardson variety X_y^x where $y = w_{max}(i_1 \dots i_j)$ and $x = w_0 w_{max}(i_{j+1} \dots i_r)^{-1}$. Moreover, $\Gamma_{\mathbf{i}}$ is smooth and irreducible.*

Proof. The variety $\Gamma_{\mathbf{i}}$ is isomorphic to the fibred product

$$Z_{i_1 \dots i_j} \times_{F\ell(n)} Z_{i_r \dots i_{j+1}}(F_{\mathrm{op\,can}}),$$

hence

$$\begin{aligned} \mathrm{pr}_j(\Gamma_{\mathbf{i}}) &= \mathrm{pr}_j(Z_{i_1 \dots i_j}) \cap \mathrm{pr}_{r-j}(Z_{i_r \dots i_{j+1}}(F_{\mathrm{op\,can}})) \\ &= X_{w_{\max}(i_1 \dots i_j)} \cap w_0 X_{w_{\max}(i_r \dots i_{j+1})} \\ &= X_y^x. \end{aligned}$$

Eventually, we may prove that $\Gamma_{\mathbf{i}}$ is smooth and irreducible exactly as in the proof of Proposition 1.4. $\square$

Example 1.7. We consider the Richardson variety $X_w^v \subset F\ell(4)$ with $w = [4231]$ and $v = [2143]$. A flag $F = (F^1 \subset F^2 \subset F^3 \subset F^4 = k^4)$ belongs to the Schubert variety X_w if and only if F^2 meets $\langle e_1, e_2 \rangle$.

Since $w = s_1 s_2 s_3 s_2 s_1$ is a reduced decomposition, the Bott-Samelson variety Z_{12321} desingularizes X_w. An element of Z_{12321} is a gallery

$$F_{\mathrm{can}} \overset{1}{\text{—}} F_1 \overset{2}{\text{—}} F_2 \overset{3}{\text{—}} F_3 \overset{2}{\text{—}} F_4 \overset{1}{\text{—}} F_5.$$

A flag G belongs to the opposite Schubert variety X^v if and only if $G^1 \subset \langle e_2, e_3, e_4 \rangle$ and $G^3 \supset \langle e_4 \rangle$.

Similarly, $w_0 v = s_2 s_1 s_3 s_2$ is a reduced decomposition, so the Bott-Samelson variety $Z_{2312}(F_{\mathrm{op\,can}})$ desingularizes the opposite Schubert variety X^v. An element of $Z_{2132}(F_{\mathrm{op\,can}})$ is a gallery

$$F_{\mathrm{op\,can}} \overset{2}{\text{—}} G_8 \overset{1}{\text{—}} G_7 \overset{3}{\text{—}} G_6 \overset{2}{\text{—}} G_5.$$

Therefore, an element of the variety $\Gamma_{123212312}$ has the form

$$\gamma = \big(F_{\mathrm{can}} \overset{1}{\text{—}} F_1 \overset{2}{\text{—}} F_2 \overset{3}{\text{—}} F_3 \overset{2}{\text{—}} F_4 \overset{1}{\text{—}} F_5 \overset{2}{\text{—}} G_6 \overset{3}{\text{—}} G_7 \overset{1}{\text{—}} G_8 \overset{2}{\text{—}} F_{\mathrm{op\,can}}\big).$$

The projection

$$\mathrm{pr}_5 : \gamma \mapsto F_5 = G_5$$

maps $\Gamma_{123212312}$ birationally to X_w^v.

There are only two singular points on X_w^v, namely e_w and e_v. Their fibres $\mathrm{pr}_5^{-1}(e_w)$ and $\mathrm{pr}_5^{-1}(e_v)$ are 1-dimensional. Indeed, given a gallery $\gamma \in \Gamma_{\mathbf{i}}$, let V_j be the i_j-th component of $\mathrm{pr}_j(\gamma)$. Since $\mathrm{pr}_{j-1}(\gamma) \overset{i_j}{\text{—}} \mathrm{pr}_j(\gamma)$, we know $\mathrm{pr}_j(\gamma)$ as soon as we know $\mathrm{pr}_{j-1}(\gamma)$ and V_j. Thus, a gallery can be given by the sequence $V_1, \ldots, V_9$. With this description, a gallery in the fibre of e_w is then given by

$$\langle e_2 \rangle,\ \langle e_2, e_3 \rangle,\ \langle e_2, e_3, e_4 \rangle,\ \langle e_2, e_4 \rangle,\ \langle e_4 \rangle,\ \langle e_4, xe_2 + ye_3 \rangle,\ \langle e_2, e_3, e_4 \rangle,\ \langle e_4 \rangle,\ \langle e_3, e_4 \rangle,$$

with $[x : y] \in \mathbf{P}^1$.

Similarly, the fibre of e_v is given by

$$\langle xe_1 + ye_2 \rangle,\ \langle e_1, e_2 \rangle,\ \langle e_1, e_2, e_3 \rangle,\ \langle e_1, e_2 \rangle,\ \langle e_2 \rangle,\ \langle e_2, e_4 \rangle,\ \langle e_2, e_3, e_4 \rangle,\ \langle e_4 \rangle,\ \langle e_3, e_4 \rangle,$$

with $[x : y] \in \mathbf{P}^1$.

2. Background on SMT for Bott-Samelson varieties

In this section, we recall from [28] the main definitions and results about Standard Monomial Theory for Bott-Samelson varieties.

Definitions 2.1. A *tableau* is a sequence $T = t_1 \ldots t_p$ with $t_j \in [n]$. If $T = t_1 \ldots t_p$ and $T' = t'_1 \ldots t'_{p'}$ are two tableaux, then the *concatenation* $T * T'$ is the tableau $t_1 \ldots t_p\, t'_1 \ldots t'_{p'}$. We denote by $\emptyset$ the empty tableau, so that $T * \emptyset = \emptyset * T = T$.

A *column κ of size i* is a tableau $\kappa = t_1 \ldots t_i$ with $1 \leq t_1 < \cdots < t_i \leq n$. The set of all columns of size i is denoted by $I_{i,n}$. The *Bruhat order* on $I_{i,n}$ is defined by

$$\kappa = t_1 \ldots t_i \leq \kappa' = t'_1 \ldots t'_i \iff t_1 \leq t'_1,\ \ldots,\ t_i \leq t'_i.$$

The symmetric group S_n acts on $I_{i,n}$: if $w \in S_n$ and $\kappa = t_1 \dots t_i \in I_{i,n}$, then $w\kappa$ is the column obtained by rearranging the tableau $w(t_1) \dots w(t_i)$ in an increasing sequence.

For $i \in [n]$, the *fundamental weight column* ϖ_i is the sequence $12 \dots i$.

We shall be interested in a particular type of tableau, called standard.

Definitions 2.2. Let $\mathbf{i} = i_1 \dots i_r$, and $\mathbf{m} = m_1 \dots m_r \in \mathbf{Z}^r_{\geq 0}$. A *tableau of shape* $(\mathbf{i}, \mathbf{m})$ is a tableau of the form

$$\kappa_{11} * \dots * \kappa_{1m_1} * \kappa_{21} * \dots * \kappa_{2m_2} * \dots * \kappa_{r1} * \dots * \kappa_{rm_r}$$

where κ_{km} is a column of size i_k for every k, m. (If $m_k = 0$, there is no column in the corresponding position of T.)

A *lifting* of T is a sequence of subwords of $\mathbf{i}$

$$J_{11} \supset \dots \supset J_{1m_1} \supset J_{21} \supset \dots \supset J_{2m_2} \supset \dots \supset J_{r1} \supset \dots \supset J_{rm_r}$$

such that $J_{km} \cap [k]$ is a reduced subword of $\mathbf{i}$ and $w(\mathbf{i}(J_{km} \cap [k]))\varpi_{i_k} = \kappa_{km}$. If such a lifting exists, then the tableau T is said to be *standard*.

Remark 2.3. The last equality in the definition of a lifting may be viewed geometrically as follows. If $J \subset [r]$ and $j \in [r]$, then $\mathrm{pr}_j : Z_{\mathbf{i}} \to F\ell(n)$ maps $Z_J \subset Z_{\mathbf{i}}$ onto a Schubert variety $X_w \subset F\ell(n)$ (*cf.* Proof of Proposition 1.3). In the notations of Section 1, the images of T-fixed points of Z_J under pr_j are of the form $\mathrm{pr}_j(e_K) = e_{u_1 \dots u_j} = e_{w(\mathbf{i}(K \cap [j]))}$ with K running over all subsets of J, hence $w = w_{\max}(\mathbf{i}(J \cap [j]))$. In turn, the image of $\mathrm{pr}_j(Z_J)$ by the projection $F\ell(n) \to G_{i_j,n}$ is equal to the Schubert variety $X_{w\varpi_{i_j}}$: for $J = J_{km}$ in the above lifting, this projection is therefore equal to $X_{\kappa_{km}}$. We shall follow up on this point of view in Remark 4.7.

Notation 2.4. Each column $\kappa \in I_{i,n}$ identifies with a weight of $GL(n)$, in such a way that the fundamental weight column ϖ_i corresponds to the ith fundamental weight of $GL(n)$. Therefore, we also denote by ϖ_i this fundamental weight.

We recall the Plücker embedding: given an i-subspace V of k^n, choose a basis $v_1, \dots, v_i$ of V, and let M be the matrix of the vectors $v_1, \dots, v_i$ written in the basis $(e_1, \dots, e_n)$. We associate to each column $\kappa = t_1 \dots t_i$ the minor $p_\kappa(V)$ of M on rows $t_1, \dots, t_i$. Then the map $p : V \mapsto [p_\kappa(V) \mid \kappa \in I_{i,n}]$ is the Plücker embedding.

Let $\pi_i : F\ell(n) \to G_{i,n}$ be the natural projection. We denote by L_{ϖ_i} the line bundle $(p \circ \pi_i)^* \mathcal{O}(1)$.

Now consider the tensor product $L_{\varpi_{i_1}}^{\otimes m_1} \otimes \dots \otimes L_{\varpi_{i_r}}^{\otimes m_r}$ on $F\ell(n)^r$, and denote by $L_{\mathbf{i},\mathbf{m}}$ its restriction to $Z_{\mathbf{i}} \subset F\ell(n)^r$.

Definition 2.5. To a tableau $T = \kappa_{11} * \dots * \kappa_{1m_1} * \dots * \kappa_{r1} * \dots * k_{rm_r}$, one associates the section $p_T = p_{\kappa_{11}} \otimes \dots \otimes p_{\kappa_{1m_1}} \otimes \dots \otimes p_{\kappa_{r1}} \otimes \dots \otimes p_{\kappa_{rm_r}}$ of $L_{\mathbf{i},\mathbf{m}}$. If T is standard of shape $(\mathbf{i}, \mathbf{m})$, then p_T is called a *standard monomial of shape* $(\mathbf{i}, \mathbf{m})$.

Theorem 2.6 ([28])**.**

1. *The standard monomials of shape* $(\mathbf{i}, \mathbf{m})$ *form a basis of the space of sections* $H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$.
2. *For* $i > 0$, $H^i(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) = 0$.
3. *The variety* $Z_{\mathbf{i}}$ *is projectively normal for any embedding induced by a very ample line bundle* $L_{\mathbf{i},\mathbf{m}}$.

3. Linear Independence

Example 3.1. We want to see on Example 1.7 how one may construct an SMT for the varieties $\Gamma_{\mathbf{i}}$.

Consider the line bundle $L_{\mathbf{i},\mathbf{m}}$ on $Z_{\mathbf{i}}$ where $\mathbf{i} = 123212312$ and $\mathbf{m} = 200010111$. We consider the restriction map $H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$, and a natural idea to get a basis of $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$ is to take all the standard monomials that do not belong to its kernel.

So let $T = \kappa_{11} * \kappa_{12} * \kappa_{51} * \kappa_{71} * \kappa_{81} * \kappa_{91}$ be a tableau of shape $(\mathbf{i}, \mathbf{m})$. The monomial p_T does not vanish identically on $\Gamma_{\mathbf{i}}$ if and only if $\kappa_{11}, \kappa_{12} \in \{1, 2\}$, $\kappa_{51} \neq 1$, $\kappa_{71} = 234$, $\kappa_{81} = 4$, $\kappa_{91} = 34$.

One may check (by computer) that there are 708 standard tableaux. Among these tableaux, 9 do not vanish identically:

$$\begin{array}{ll}
T_1 = 2*2*\emptyset*\emptyset*\emptyset*4*\emptyset*234*4*34 & T_4 = 2*1*\emptyset*\emptyset*\emptyset*4*\emptyset*234*4*34 \\
T_2 = 2*2*\emptyset*\emptyset*\emptyset*3*\emptyset*234*4*34 & T_5 = 2*1*\emptyset*\emptyset*\emptyset*3*\emptyset*234*4*34 \\
T_3 = 2*2*\emptyset*\emptyset*\emptyset*2*\emptyset*234*4*34 & T_6 = 2*1*\emptyset*\emptyset*\emptyset*2*\emptyset*234*4*34
\end{array}$$

$$\begin{array}{l}
T_7 = 1*1*\emptyset*\emptyset*\emptyset*4*\emptyset*234*4*34 \\
T_8 = 1*1*\emptyset*\emptyset*\emptyset*3*\emptyset*234*4*34 \\
T_9 = 1*1*\emptyset*\emptyset*\emptyset*2*\emptyset*234*4*34
\end{array}$$

Moreover, the tableaux T_i admit the following liftings (J^i_{km})

$$\begin{array}{lll}
J^1_{11} = \{1,2,3,4,5,6,7,8,9\} & J^4_{11} = \{1,2,3,4,5,6,7,8,9\} & J^7_{11} = \{2,3,4,5,6,7,8,9\} \\
J^1_{21} = \{1,2,3,4,5,6,7,8,9\} & J^4_{21} = \{\quad 2,3,4,5,6,7,8,9\} & J^7_{21} = \{2,3,4,5,6,7,8,9\} \\
J^1_{51} = \{\qquad 3,4,5,6,7,8,9\} & J^4_{51} = \{\quad 2,3,4,5,6,7,8,9\} & J^7_{51} = \{2,3,4,5,6,7,8,9\} \\
J^1_{71} = \{\qquad 3,4,5,6,7,8,9\} & J^4_{71} = \{\quad 2,3,4,5,6,7,8,9\} & J^7_{71} = \{2,3,4,5,6,7,8,9\} \\
J^1_{81} = \{\qquad 3,4,5,6,7,\quad 9\} & J^4_{81} = \{\quad 2,3,4,5,6,7,\quad 9\} & J^7_{81} = \{2,3,4,5,6,7,\quad 9\} \\
J^1_{91} = \{\qquad 3,4,5,6,7,\quad 9\} & J^4_{91} = \{\quad 2,3,4,5,6,7\qquad\} & J^7_{91} = \{2,3,4,5,6,7\qquad\}
\end{array}$$

$$\begin{array}{lll}
J^2_{11} = \{1,2,3,4,5,6,7,8,9\} & J^5_{11} = \{1,2,3,4,5,6,7,8,9\} & J^8_{11} = \{2,3,4,5,6,7,8,9\} \\
J^2_{21} = \{1,2,3,4,5,6,7,8,9\} & J^5_{21} = \{\quad 2,3,4,5,6,7,8,9\} & J^8_{21} = \{2,3,4,5,6,7,8,9\} \\
J^2_{51} = \{1,2,3,\quad 5,6,7,8,9\} & J^5_{51} = \{\quad 2,3,\quad 5,6,7,8,9\} & J^8_{51} = \{2,3,\quad 5,6,7,8,9\} \\
J^2_{71} = \{1,2,3,\quad 5,6,\quad 8,9\} & J^5_{71} = \{\quad 2,3,\quad 5,6,7,8,9\} & J^8_{71} = \{2,3,\quad 5,6,7,8,9\} \\
J^2_{81} = \{1,2,3,\quad 5,6,\quad 8,9\} & J^5_{81} = \{\quad 2,3,\quad 5,6,7,8,9\} & J^8_{81} = \{2,3,\quad 5,6,7,8,9\} \\
J^2_{91} = \{1,2,3,\quad 5,6,\quad 8\quad\} & J^5_{91} = \{\quad 2,3,\quad 5,6,7,8\quad\} & J^8_{91} = \{2,3,\quad 5,6,7,8\quad\}
\end{array}$$

$$\begin{array}{lll}
J^3_{11} = \{1,2,3,4,5,6,7,8,9\} & J^6_{11} = \{1,2,3,4,5,6,7,8,9\} & J^9_{11} = \{2,3,4,5,6,7,8,9\} \\
J^3_{21} = \{1,2,3,4,5,6,7,8,9\} & J^6_{21} = \{\quad 2,3,4,5,6,7,8,9\} & J^9_{21} = \{2,3,4,5,6,7,8,9\} \\
J^3_{51} = \{1,2,3,4,\quad 6,7,8,9\} & J^6_{51} = \{\qquad 3,\quad 5,6,7,8,9\} & J^9_{51} = \{\quad 3,\quad 5,6,7,8,9\} \\
J^3_{71} = \{1,2,3,4,\qquad\quad 8,9\} & J^6_{71} = \{\qquad 3,\quad 5,6,7,8,9\} & J^9_{71} = \{\quad 3,\quad 5,6,7,8,9\} \\
J^3_{81} = \{1,2,3,4,\qquad\quad 8,9\} & J^6_{81} = \{\qquad 3,\quad 5,6,7,8,9\} & J^9_{81} = \{\quad 3,\quad 5,6,7,8,9\} \\
J^3_{91} = \{1,2,3,4,\qquad\quad 8,9\} & J^6_{91} = \{\qquad 3,\quad 5,6,7,8,9\} & J^9_{91} = \{\quad 3,\quad 5,6,7,8,9\}
\end{array}$$

These liftings have the following property: $w_{\max}(\mathbf{i}(J^i_{km})) = w_0$ for each k, m. We then say that T_i is w_0-standard. It can be checked that the standard tableaux that are not w_0-standard vanish identically on $\Gamma_{\mathbf{i}}$.

To see that the remaining monomials p_{T_i} are linearly independent, we may work on an open affine set. There exists an open set Ω of $Z_{\mathbf{i}}$ such that $\Gamma_{\mathbf{i}} \cap \Omega$ is isomorphic to the affine space k^3 (see Definition 5.10 and Proposition 5.19). Here, we have

$$\varphi : (x, y, z) \mapsto (V_1, \dots, V_9),$$

for

$$\begin{array}{ll}
V_1 = \langle xe_1 + e_2 \rangle & V_2 = \langle xe_1 + e_2, -xye_1 + e_3 \rangle \\
V_3 = \langle xe_1 + e_2, -xye_1 + e_3, e_4 \rangle & V_4 = \langle xe_1 + e_2, -xyze_1 + ze_3 + e_4 \rangle \\
V_5 = \langle yze_2 + ze_3 + e_4 \rangle & V_6 = \langle yze_2 + ze_3 + e_4, ye_2 + e_3 \rangle \\
V_7 = \langle e_2, e_3, e_4 \rangle & V_8 = \langle e_4 \rangle \\
V_9 = \langle e_3, e_4 \rangle &
\end{array}$$

We denote again by p_T the polynomial $\varphi^* \left((p_T)_{|\Omega} \right)$. We then have

$$\begin{array}{lll}
p_{T_1} = 1, & p_{T_4} = x, & p_{T_7} = x^2, \\
p_{T_2} = z, & p_{T_5} = xz, & p_{T_8} = x^2 z, \\
p_{T_3} = yz, & p_{T_6} = xyz, & p_{T_9} = x^2 yz.
\end{array}$$

It is clear that these monomials are linearly independent in $k[x, y, z]$.

Definitions 3.2 below will generalize the behaviour of the liftings (J^i_{km}) observed in this example.

Definitions 3.2. Let T be a standard tableau of shape $(\mathbf{i}, \mathbf{m})$. We say that T (or the monomial p_T) is *w_0-standard* if there exists a lifting (J_{km}) of T such that each subword J_{km} contains a reduced expression of w_0.

More generally, if $J \subset [r]$ contains a reduced expression for w_0, then $\Gamma_J = Z_J \cap \Gamma_{\mathbf{i}} \neq \emptyset$, and we say that T (or p_T) is *w_0-standard on Γ_J* if there exists a lifting (J_{km}) of T such that for every k, m, $J \supset J_{km}$ and J_{km} contains a reduced expression of w_0.

Similarly, T (or p_T) is said to be *w_0-standard on a union* $\Gamma = \Gamma_{J_1} \cup \cdots \cup \Gamma_{J_k}$ if T is w_0-standard on at least one of the components $\Gamma_{J_1}, \dots, \Gamma_{J_k}$. We then denote by $\mathcal{S}(\Gamma)$ the set of all w_0-standard tableaux on Γ.

We need some results about *positroid varieties.* References for these varieties can be found in [21].

Let π_i be the canonical projection $F\ell(n) \to G_{i,n}$. In general, the projection of a Richardson variety $X^v_w \subset F\ell(n)$ is no longer a Richardson variety. But $\pi_i(X^v_w)$ is still defined inside the Grassmannian $G_{i,n}$ by the vanishing of some Plücker coordinates. More precisely, consider the set $\mathcal{M} = \{\kappa \in I_{i,n} \mid e_\kappa \in \pi_i(X^v_w)\}$. Then

$$\Pi = \pi_i(X^v_w) = \{V \in G_{i,n} \mid \kappa \notin \mathcal{M} \implies p_\kappa(V) = 0\}.$$

The poset $\mathcal{M}$ is a *positroid* (see the paragraph following Lemma 3.20 in [21]), and the variety Π is called a *positroid variety.*

Lemma 3.3. *With the notation above,*

$$\mathcal{M} = \{\kappa \in I_{i,n} \mid \exists u \in [v, w],\ u\varpi_i = \kappa\}.$$

Proof. Let $u \in [v, w]$ and $\kappa = u\varpi_i$. Then $e_u \in X_w^v$, so $e_\kappa = \pi_i(e_u) \in \Pi$. Hence $\kappa \in \mathcal{M}$.

Conversely, let $\kappa \in \mathcal{M}$. The fibre $\pi_i^{-1}\{e_\kappa\}$ in X_w^v is a non-empty T-stable variety, hence, by Borel's fixed point theorem, this variety has a T-fixed point e_u, $u \in S_n$. It follows that $u \in [v, w]$ and $u\varpi_i = \kappa$. □

Theorem 3.4. *For every subword $J_1, \ldots, J_k$ containing a reduced expression of w_0, the w_0-standard monomials on the union $\Gamma = \Gamma_{J_1} \cup \cdots \cup \Gamma_{J_k}$ are linearly independent.*

Proof. We imitate the proof of the corresponding proposition for Bott-Samelson varieties appearing in [28, Section 3.2]. Let $\mathcal{T}$ be a non-empty subset of $\mathcal{S}(\Gamma)$, and assume that we are given a linear relation among monomials p_T for T in $\mathcal{T}$:

$$(*) \qquad \sum_{T \in \mathcal{T}} a_T \, p_T(\gamma) = 0 \quad \forall \gamma \in \Gamma.$$

Moreover, we may assume that the coefficients appearing in this relation are all non-zero. We shall proceed by induction on the length of tableaux, that is, on $M = \sum_{i=1}^r m_i$.

If $M = 1$, then $\mathbf{m}$ has the form $0 \ldots 1 \ldots 0$, that is, we have $m_e = 1$ for some e, and $m_i = 0$ for all $i \neq e$. The tableaux T that appear in relation $(*)$ are of the form $T = \kappa_e$, where $\kappa_e \in I_{i_e,n}$. If $\gamma = (F_{\mathrm{can}}, F_1, \ldots, F_r = F_{\mathrm{op\,can}}) \in \Gamma$ then $p_T(\gamma) = p_{\kappa_e}(F_e)$. Thus, we have a linear relation of Plücker coordinates in a union of Richardson varieties in $F\ell(n)$, hence a linear relation on one of these Richardson varieties. But Standard Monomial Theory for Richardson varieties (*cf.* [25], Theorem 32) shows that such a relation cannot exist.

Now assume that $M > 1$, and $\mathbf{m} = 0 \ldots 0 \, m_e \ldots m_r$ with $m_e > 0$. Here, we denote by κ_{km}^T the columns of a tableau T. Consider an element κ minimal among the first columns of the tableaux of $\mathcal{T}$, that is,

$$\kappa \in \min\{\kappa_{e1}^T \mid T \in \mathcal{T}\}.$$

We consider the set $\mathcal{T}(\kappa)$ of tableaux T in $\mathcal{T}$ with $\kappa_{e1}^T = \kappa$. For every $T \in \mathcal{T}(\kappa)$, fix a maximal lifting $J_{e1}^T \supset \cdots \supset J_{rm_r}^T$ containing a reduced expression of w_0 and with J_{e1}^T contained in one of the subwords $J_1, \ldots, J_k$, so that $\Gamma \supset \Gamma_{J_{e1}^T} \neq \emptyset$. Thus, we can restrict the relation $(*)$ on

$$\Gamma(\kappa) = \bigcup_{T \in \mathcal{T}(\kappa)} \Gamma_{J_{e1}^T}.$$

If $T \in \mathcal{T}(\kappa)$, then $T = \kappa * T'$, and T' is a w_0-standard tableau on $\Gamma(\kappa)$ of shape $(\mathbf{i}, 0 \ldots 0 \, m_e - 1 \ldots m_r)$.

If $T \notin S(\kappa)$, then $\kappa_{e1}^T \not\leq \kappa$, so $p_{\kappa_{e1}^T}$ vanishes identically on the Schubert variety $X_\kappa \subset G_{i_e,n}$, hence on each Schubert variety $X_{w_{\max}(\mathbf{i}(J_{e1}^S))}$ for $S \in \mathcal{T}(\kappa)$. In particular, $p_{\kappa_{e1}^T}$ vanishes on $\Gamma(\kappa)$, and p_T as well.

Restrict relation $(*)$ to $\Gamma(\kappa)$:

$$p_\kappa(\gamma) \sum_{T \in \mathcal{T}(\kappa)} a_T \, p_{T'}(\gamma) = 0 \quad \forall \gamma \in \Gamma(\kappa).$$

This product vanishes on each irreducible $\Gamma_{J_{e1}^T}$ $(T \in \mathcal{T}(\kappa))$. Now, p_κ does not vanish identically on $\Gamma(J_{e1}^T)$. Indeed, we know by Proposition 1.6 that $\mathrm{pr}_e(\Gamma(J_{e1}^T))$ is the Richardson variety X_y^x with $y = w(\mathbf{i}(J_{e1}^T)) \geq x$. Since $\kappa = y\varpi_{i_e}$, by Lemma 3.3, p_κ does not vanish identically on X_y^x, hence does not vanish identically on $\Gamma(J_{e1}^T)$.

So we may simplify by p_κ on the irreducible $\Gamma_{J_{e1}^T}$, hence a linear relation between w_0-standard monomials on $\Gamma(\kappa)$ of shape $(\mathbf{i}, 0 \dots 0\, m_e - 1 \dots m_r)$. By induction over M, $a_T = 0$ for all $T \in \mathcal{T}(\kappa)$: a contradiction. □

4. Standard monomials that do not vanish on $\Gamma_{\mathbf{i}}$ are w_0-standard

In this section, we shall prove that the standard monomials that do not vanish identically on $\Gamma_{\mathbf{i}}$ are w_0-standard, provided certain assumptions over $\mathbf{m}$, which cover the regular case (*i.e.* $m_i > 0$ for every i).

Lemma 4.1 (Lifting Property [3, Proposition 2.2.7])**.** *Let s be a simple transposition, and $u < w$ in S_n.*

- *If $u < su$ and $w > sw$, then $u \leq sw$ and $su \leq w$.*
- *If $u > su$ and $w > sw$, then $su \leq sw$.*
- *If $u < su$ and $w < sw$, then $su \leq sw$.* □

We may represent these situations by the pictures below







Definition 4.2. Let $x, y \in S_n$. The Demazure product $x * y$ is the unique maximal element of the poset $\mathcal{D}(x, y) = \{uv \mid u \leq x, v \leq y\}$.

Proposition 4.3 ([37, 18])**.** *Let $x, y \in S_n$. The double coset $B(x*y)B$ is the unique $B \times B$-double coset that is open in $BxByB$. In particular, $*$ is associative.* □

Lemma 4.4. *Let s be a simple transposition, and $x \in S_n$. Then $x * s = \max(x, xs)$. Similarly, $s * x = \max(x, sx)$.*

Proof. We shall prove that $x * s = \max(x, xs)$, the proof of $s * x = \max(x, sx)$ being similar.

- Case 1: $x > xs$. Let $u \leq x$. If $us < u$, then $us \leq x$. If $us > u$, then by Lemma 4.1, we have $us \leq x$. Hence every element of $\mathcal{D}(x, s)$ is less than or equal to x, so $x * s = x = \max(x, xs)$.
- Case 2: $x < xs$. Let $u \leq x$. If $us < u$, then $us \leq xs$. If $us > u$, then by Lemma 4.1, $us \leq xs$. Thus, every element of $\mathcal{D}(x, s)$ is less than or equal to xs, so $x * s = xs = \max(x, xs)$. □

Lemma 4.5. *Let J be a subword of $\mathbf{i}$. For every $k \in [r]$,*

$$w_{max}\big(\mathbf{i}(J)\big) = w_{max}\big(\mathbf{i}(J \cap [k])\big) * w_{max}\big(\mathbf{i}(J \cap [k+1, r])\big).$$

Proof. Let

$$\begin{aligned} w =& w_{\max}\big(\mathbf{i}(J)\big) \\ x =& w_{\max}\big(\mathbf{i}(J \cap [k])\big) \\ y =& w_{\max}\big(\mathbf{i}(J \cap [k+1, r])\big) \end{aligned}$$

Each element uv of $\mathcal{D}(x, y)$ has a decomposition of the form $w\big(\mathbf{i}(K_1)\big)w\big(\mathbf{i}(K_2)\big)$ with $K_1 \subset J \cap [k]$ and $K_2 \subset J \cap [k+1, r]$. Hence,

$$uv = w\big(\mathbf{i}(K_1 \cup K_2)\big) \leq w,$$

so $x * y \leq w$.

Conversely, let $K' \subset J$ be such that $w\big(\mathbf{i}(K')\big) = w$ is a reduced decomposition. Since

$$w = w\big(\mathbf{i}(K' \cap [k])\big)w\big(\mathbf{i}(K' \cap [k+1, r])\big),$$

we have $w \in D(x, y)$, hence $w \leq x * y$. □

Lemma 4.6 ([17, 2.2.(4)])**.** *If $x' \leq x$ and $y' \leq y$, then $x' * y' \leq x * y$.*

Proof. By Proposition 4.3, we have

$$B(x' * y')B \subset \overline{BxB}\ \overline{ByB} \subset \overline{BxByB} = \overline{B(x * y)B},$$

hence $x' * y' \leq x * y$. □

Let T be a standard tableau of shape $(\mathbf{i}, \mathbf{m})$, and e be the least integer such that $m_e \neq 0$, so $\mathbf{m} = 0 \dots 0\, m_e \dots m_r$. We give the construction of a particular type of liftings of T (called optimal), in light of the following

Remark 4.7. Let (K_{km}) be an arbitrary lifting of T and set

$$w_{km} = w\big(\mathbf{i}(K_{km} \cap [k])\big),$$

so that $w_{km}\varpi_k = \kappa_{km}$. By Remark 2.3, $\mathrm{pr}_k(Z_{K_{km}}) = X_{w_{km}}$, with the following consequences.

- For each k, $K_{k1} \supset \cdots \supset K_{km_k}$ yields $w_{k1} \geq \cdots \geq w_{km_k}$.
- Let l be the least integer such that $l > k$ and $m_l \neq 0$. Then $K_{km_k} \supset K_{l,1}$ yields $\mathrm{pr}_l(Z_{K_{l,1}}) \subset \mathrm{pr}_l(Z_{K_{km_k}})$, hence

  $$w\big(\mathbf{i}(K_{l,1} \cap [l])\big) \leq w_{\max}\big(\mathbf{i}(K_{km_k} \cap [l])\big).$$

  By Lemma 4.5,

  $$w_{\max}\big(\mathbf{i}(K_{km_k} \cap [l])\big) = w\big(\mathbf{i}(K_{k,m_k} \cap [k])\big) * w_{\max}\big(\mathbf{i}(K_{k,m_k} \cap [k+1, l])\big).$$

  So

  $$w_{l,1} \leq w_{k,m_k} * w_{\max}\big(\mathbf{i}(K_{k,m_k} \cap [k+1, l])\big).$$

We shall also need a result due to V. Deodhar:

Notation 4.8. Let $\kappa \in I_{i,n}$ and $w \in S_n$. We set

$$\mathcal{E}(w, \kappa) = \{v \in S_n \mid v \leq w,\ v\varpi_i = \kappa\}.$$

Lemma 4.9 ([27, Lemma 11])**.** *Let $\kappa \in I_{i,n}$, and $w \in S_n$. If $\mathcal{E}(w, \kappa) \neq \emptyset$, then it admits a unique maximal element.* □

Remark 4.10. The above lemma admits the following geometric interpretation. Let q be the restriction to X_w of the canonical projection $F\ell(n) \to G_{i,n}$. Since q is B-equivariant, $q^{-1}(X_\kappa)$ is a union of Schubert varieties, namely

$$q^{-1}(X_\kappa) = \bigcup_{v \in \mathcal{E}(x,\kappa)} X_v.$$

By Lemma 4.9, we conclude that $q^{-1}(X_\kappa)$ is a single Schubert variety.

There exists a dual version of the lemma: if $w\varpi_i \geq \kappa$, then the set

$$\{v \in S_n \mid v \geq w,\ v\varpi_i = \kappa\}$$

admits a unique minimal element. Hence, $q^{-1}(X^\kappa)$ is a single opposite Schubert variety.

We now construct elements $v_{km} \in S_n$ inductively, as follows. At the first step, consider the set

$$\mathcal{E}(w_{\max}(i_1 \dots i_e), \kappa_{e1}).$$

Since it contains w_{e1}, it is nonempty, so it has a maximal element v_{e1}, which is unique thanks to Lemma 4.9. Now assume that $v_{km} \geq w_{km}$ has already been constructed. We then proceed in the same way to construct $v_{k,m+1}$ (if $m < m_k$) or $v_{l,1}$ (if $m = m_k$, and $l > k$ is the least integer such that $m_l \neq 0$):

- If $m < m_k$, then the set $\mathcal{E}(v_{k,m}, \kappa_{k,m+1})$ is nonempty (since it contains $w_{k,m+1}$), so let $v_{k,m+1}$ be its unique maximal element.
- If $m = m_k$, then let $v'_{k,m} = v_{k,m} * w_{\max}\big(i_{k+1} \dots i_l\big)$. By Lemma 4.6,

  $$w_{k,m} * w_{\max}\big(\mathbf{i}(K_{k,m} \cap [k+1, l])\big) \leq v'_{k,m}.$$

  Thus, by Remark 4.7 the set $\mathcal{E}(v'_{k,m}, \kappa_{l,1})$ contains $w_{l,1}$, so it is non-empty. Let $v_{l,1}$ be its unique maximal element.

Remark 4.11. Although the *existence* of the v_{km} depends on that of the w_{km} (*i.e.* on the existence of a lifting of the tableau T), the *values* of the v_{km} only depend on the tableau T itself.

Next, we construct subsets $E_{km} \subset [k]$, again inductively. Since

$$v_{e1} \leq w_{\max}(i_1 \dots i_e),$$

choose $E_{e1} \subset \{i_1 \dots i_e\}$ such that v_{e1} admits a reduced expression of the form $\mathbf{i}(E_{e1})$. If $E_{k,m}$ such that $v_{k,m} = w\big(\mathbf{i}(E_{k,m})\big)$ has already been constructed, then define $E_{k,m+1}$ (if $m < m_k$) or $E_{l,1}$ (if $m = m_k$) as follows:

- If $m < m_k$, then $v_{k,m+1} \leq v_{k,m} = w\big(\mathbf{i}(E_{k,m})\big)$, so choose $E_{k,m+1} \subset E_{k,m}$ such that $v_{k,m+1}$ admits a reduced expression of the form $\mathbf{i}(E_{k,m+1})$.
- If $m = m_k$, then by Lemma 4.5

  $$v_{l,1} \leq v'_{k,m_k} = w_{\max}\big(\mathbf{i}(E_{km_k} \cup \{k+1, \dots, l\})\big),$$

  so choose $E_{l,1} \subset E_{km_k} \cup \{k+1, \dots, l\}$ such that $v_{l,1}$ admits a reduced expression of the form $\mathbf{i}(E_{l,1})$.

Definition 4.12. With the above notation, set $J_{km} = E_{km} \cup [k+1, r]$ for each k, m. We will call (J_{km}) an *optimal lifting* of the tableau T.

Remark 4.13. The optimal lifting is not unique. However, while it depends on the choice of reduced expressions for the v_{km}, it is still independent on the choice of the initial lifting (K_{km}).

Example 4.14. Consider the tableau $T = 123 * 13 * 3 * 134 * 24 * 124$ of shape $(3213233213, 1111110000)$. This tableau is standard, and we shall construct an optimal lifting of T.

- $v_{11} = \max \mathcal{E}(s_3, 123) = e$.
- $v_{21} = \max \mathcal{E}(e * s_2, 13) = s_2$.
- $v_{31} = \max \mathcal{E}(s_2 * s_1, 3) = s_2 s_1$.
- $v_{41} = \max \mathcal{E}(s_2 s_1 * s_3, 134) = s_2 s_1 s_3$.
- $v_{51} = \max \mathcal{E}(s_2 s_1 s_3 * s_2, 24) = s_1 s_3 s_2$.
- $v_{61} = \max \mathcal{E}(s_1 s_3 s_2 * s_3, 124) = s_1 s_3$.

Hence

$$\begin{aligned}
J_{11} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{21} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{31} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{41} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{51} &= \{3,4,5,6,7,8,9,10\}\\
J_{61} &= \{3,4,7,8,9,10\}
\end{aligned}$$

is an optimal lifting of T. Another optimal lifting of T is

$$\begin{aligned}
J'_{11} &= \{2,3,4,5,6,7,8,9,10\}\\
J'_{21} &= \{2,3,4,5,6,7,8,9,10\}\\
J'_{31} &= \{2,3,4,5,6,7,8,9,10\}\\
J'_{41} &= \{2,3,4,5,6,7,8,9,10\}\\
J'_{51} &= \{3,4,5,6,7,8,9,10\}\\
J'_{61} &= \{3,6,7,8,9,10\}
\end{aligned}$$

Lemma 4.15. *Let $w \in S_n$ and $\kappa \in I_{i,n}$ be such that $\mathcal{E}(w, \kappa) \neq \emptyset$. Consider a simple transposition s such that $sw < w$.*

(1) If $s\kappa > \kappa$, then $\max \mathcal{E}(w, \kappa) = \max \mathcal{E}(sw, \kappa)$.
(2) If $s\kappa \leq \kappa$, then $\max \mathcal{E}(w, \kappa) = s * \max \mathcal{E}(sw, s\kappa)$.

Proof. Let $u = \max \mathcal{E}(w, \kappa)$.

- Case 1: assume that $su > u$. Then, by Lemma 4.1,



we have $u \leq sw$ and $su \leq w$. Hence $s\kappa \geq \kappa$, but by maximality of u, $su \notin \mathcal{E}(w,\kappa)$, hence $s\kappa > \kappa$. Since $u \leq sw$, $u \in \mathcal{E}(sw,\kappa)$, so

$$u \leq \max \mathcal{E}(sw,\kappa) \leq \max \mathcal{E}(w,\kappa) = u.$$

This proves the part (1) of the lemma.

- Case 2: $su < u$. Then $s\kappa \leq \kappa$, and by Lemma 4.1,



we have $su \leq sw$, so $su \in \mathcal{E}(sw, s\kappa)$, hence $su \leq v = \max \mathcal{E}(sw, s\kappa)$. We distinguish two subcases:
  - Subcase 1: $s\kappa < \kappa$. Then $sv > v$. Since we also have $su < u$, it follows from Lemma 4.1 that $v \leq su$. Similarly, $sv > v$, together with $sw < w$ imply that $sv \leq w$, so $sv \in \mathcal{E}(w,\kappa)$, hence $sv \leq u$. By Lemma 4.1, we have $v \leq su$. So $v = su$, or equivalently

$$u = sv = \max(v, sv) = s * v.$$

  - Subcase 2: $s\kappa = \kappa$.
    - ∗ If $u \leq sw$, then $u \in \mathcal{E}(sw,\kappa)$, so $u \leq v$. But $v \leq u$, so $u = v$.
    - ∗ If $u \nleq sw$, then $su \in \mathcal{E}(sw,\kappa)$, so $su \leq v \leq u$. In other words, $v \in \{u, su\}$.

  In each of these two situations, we have $u = v$ or $u = sv$. But, if $sv > v$ then $u \neq v$ (since $su < u$), so $u = sv = \max(v, sv) = s * v$. If $sv < v$, then $u \neq sv$, so

$$u = v = s * v. \ \square$$

Let $w = s_{i_1} \dots s_{i_j}$ be a reduced expression. The lemma above gives an algorithm to find a reduced expression of $u = \max \mathcal{E}(w,\kappa)$, say $u = w(\mathbf{i}(J))$, with $J \subset [j]$: let $s = s_{i_1}$, and compare $s\kappa$ with κ.

- If $s\kappa > \kappa$, then $u = \max \mathcal{E}(sw,\kappa)$.
- If $s\kappa \leq \kappa$, then $u = s * \max \mathcal{E}(sw, s\kappa)$.

We then compute $\max \mathcal{E}(sw, s\kappa)$ or $\max \mathcal{E}(sw,\kappa)$ in the same way, using the decomposition $sw = s_{i_2} \dots s_{i_j}$.

Example 4.16. In S_4, take $w = [4231] = s_1 s_2 s_3 s_2 s_1$ and $\kappa = 13$. We shall compute $u = \max \mathcal{E}(w,\kappa)$ with the previous algorithm. Note that $\kappa \leq 24 = w\varpi_2$, hence $\mathcal{E}(w,\kappa) \neq \emptyset$.

- $s_1\kappa = 23 > \kappa$, so $u = \max \mathcal{E}(s_2 s_3 s_2 s_1, \kappa)$,
- $s_2\kappa = 12 \leq k$, so $u = s_2 * \max \mathcal{E}(s_3 s_2 s_1, 12)$,
- $s_3(12) = 12$, so $u = s_2 * s_3 * \max \mathcal{E}(s_2 s_1, 12)$,
- $s_2(12) = 13 > 12$, so $u = s_2 * s_3 * \max \mathcal{E}(s_1, 12)$,
- $s_1(12) = 12$, so $u = s_2 * s_3 * s_1 * \max \mathcal{E}(e, 12)$.

Now, $\max(e, 12) = e$, so $u = s_2 * s_3 * s_1 = s_2 s_3 s_1 = [3142]$.

Lemma 4.17 ([3, Proposition 2.4.4])**.** *Let $\kappa \in I_{i,n}$. The set $\{v \in S_n \mid v\varpi_i = \kappa\}$ admits a unique minimal element u. Moreover, if $v \in S_n$ satisfies $v\varpi_i = \kappa$, then v admits a unique factorization $v = uv'$ with $v'\varpi_i = \varpi_i$. This factorization is length-additive, in the sense that $l(v) = l(u) + l(v')$.* □

Remark 4.18. Lemmas 4.1, 4.4, 4.6, Definition 4.2 and Proposition 4.3 are true for every Weyl group. Lemmas 4.9, 4.17, and Remark 4.10 are also true if we replace S_n by a Weyl group W, $I_{i,n}$ by the set W^P of minimal representatives of the quotient W/W_P, W_P a parabolic subgroup of W, and $v\varpi_i = \kappa$ by $v \equiv \kappa \mod W_P$.

Lemma 4.19. *Denote by u_d the minimal permutation such that $u_d\varpi_d = w_0\varpi_d$. Let $w \geq u$, and κ a column of arbitrary size $i \leq n$ such that $\mathcal{E}(w, \kappa) \neq \emptyset$. Assume that $x = \max \mathcal{E}(w, \kappa) \geq u$. Then*

$$\forall v \geq u,\ \mathcal{E}(v, \kappa) \neq \emptyset \implies \max \mathcal{E}(v, \kappa) \geq u.$$

Proof. Since $v \geq u$, we have $v\varpi_d = w_0\varpi_d$, hence by Lemma 4.17, $v = uv'$ with v' in the stabilizer of ϖ_d. Moreover, this decomposition is length-additive, so if $u = s_{i_1} \dots s_{i_j}$ and $v' = s_{i_{j+1}} \dots s_{i_k}$ are reduced expressions, then $s_{i_1} \dots s_{i_j} s_{i_{j+1}} \dots s_{i_k}$ is a reduced expression of v. Similarly, we decompose $x = ux'$ with $x'\varpi_d = \varpi_d$. We then obtain

$$x > s_{i_1}x > \cdots > s_{i_j} \dots s_{i_1}x = x',$$

hence

$$\kappa \geq s_{i_1}\kappa \geq \cdots \geq s_{i_j} \dots s_{i_1}\kappa.$$

Now, we apply the procedure described after Lemma 4.15 for the decomposition $v = s_{i_1} \dots s_{i_j} s_{i_{j+1}} \dots s_{i_k}$. The above inequalities show that $\max \mathcal{E}(v, \kappa)$ is of the form $s_{i_1} * \cdots * s_{i_j} * z$. But, by Lemma 4.6, we have

$$\begin{aligned} s_{i_1} * \cdots * s_{i_j} * z &\geq s_{i_1} * \cdots * s_{i_j} \\ &\geq s_{i_1} \dots s_{i_j} \\ &\geq u. \ \square \end{aligned}$$

Notation 4.20. For $k \in [n-1]$, let j_k be the greatest integer such that $i_{j_k} = k$.

Theorem 4.21. *Assume that for every k, $m_{j_k} > 0$. Then the standard monomials p_T of shape $(\mathbf{i}, \mathbf{m})$ that do not vanish identically on $\Gamma_{\mathbf{i}}$ are w_0-standard.*

Proof. Consider an optimal lifting (J_{km}) of T. Let $(F_{\text{can}}, F_1, \dots, F_r) \in \Gamma_{\mathbf{i}}$ be a gallery such that $p_T(F_{\text{can}}, F_1, \dots, F_r) \neq 0$. By definition of j_k, the flags F_{j_k} and $F_{\text{op can}}$ share the same k-subspace, which then is the T-fixed point $\langle e_n, \dots, e_{n-k+1}\rangle$. Hence, $\kappa_{j_k,1} = \cdots = \kappa_{j_k,m_{j_k}} = w_0\varpi_k$.

Arrange the integers $j_1, \dots, j_{n-1}$ in an increasing sequence: $j_{l_1} < \cdots < j_{l_{n-1}}$.

We shall prove that if $k > j_l$, then $v_{km} \geq u_l$. Since $p_T(F_{\text{can}}, F_1, \dots, F_r) \neq 0$, we have $p_{\kappa_{km}}(F_k) \neq 0$, hence $p_{\kappa_{km}}$ does not vanish identically on the Richardson variety X_w^v, where $w = w_{\max}(i_1 \dots i_k)$ and $v = w_0(w_{\max}(i_{k+1} \dots i_r))^{-1}$. This means that $p_{\kappa_{km}}$ does not vanish identically on the positroid variety $\pi(X_w^v)$, where $\pi : F\ell(n) \to G_{i_k,n}$. By Lemma 3.3, there exists $u \in [v, w]$ such that $u\varpi_{i_k} = \kappa_{km}$. It follows that the maximal element x_l of $\mathcal{E}(w, \kappa_{km})$ is greater than u. But, since $k > j_l$, a reduced expression of w_0v^{-1} consists of letters taken from $i_{k+1} \dots i_r$, where no l appears. Thus, $w_0v^{-1}\varpi_l = \varpi_l$, so $v\varpi_l = w_0\varpi_l$, that is, $v \geq u_l$. Hence $x_1 \geq u \geq v \geq u_l$. We then conclude with Lemma 4.19.

Now, we consider subwords J_{km} with $k \leq j_{l_1}$. In this case, $k \leq j_t$ ($t \geq 1$), so we have the inequalities $w\big(\mathbf{i}(J_{j_t,1} \cap [j_t])\big) \leq w_{\max}\big(\mathbf{i}(J_{km})\big)$, hence

$$w_{\max}\big(\mathbf{i}(J_{km})\big)\varpi_t \geq w\big(\mathbf{i}(J_{j_t,1} \cap [j_t])\big)\varpi_t = \kappa_{j_t,1} = w_0\varpi_t,$$

i.e. $w_{\max}\big(\mathbf{i}(J_{km})\big)\varpi_t = w_0\varpi_t$. So $w_{\max}\big(\mathbf{i}(J_{km})\big) = w_0$.

If $j_{l_t} < k \leq j_{l_{t+1}}$, then we have, in one hand, $w_{\max}\big(\mathbf{i}(J_{km})\big) \geq w\big(\mathbf{i}(J_{j_{l_p},1})\big)$ for every $p \geq t+1$, so $w_{\max}\big(\mathbf{i}(J_{km})\big)\varpi_{l_p} \geq \kappa_{l_p,1} = w_0\varpi_{l_p}$, hence $w_{\max}\big(\mathbf{i}(J_{km})\big)\varpi_{l_p} = w_0\varpi_{l_p}$. On the other hand, $w_{\max}\big(\mathbf{i}(J_{km})\big) \geq v_{km} \geq u_{l_q}$ for every $q \leq t$, hence $w_{\max}\big(\mathbf{i}(J_{km})\big)\varpi_{l_q} = w_0\varpi_{l_q}$. It follows that $w_{\max}\big(\mathbf{i}(J_{km})\big) = w_0$. $\square$

Remark 4.22. The assumption $m_{j_k} > 0$ for every k is necessary: recall the tableau $T = 123{*}13{*}3{*}134{*}24{*}124$ of Example 4.14. It is standard of shape $(3213233213, 1111110000)$, and one may check that p_T does not vanish identically on $\Gamma_{\mathbf{i}}$. However, an optimal lifting of T is given by

$$\begin{aligned}
J_{11} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{21} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{31} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{41} &= \{2,3,4,5,6,7,8,9,10\}\\
J_{51} &= \{\ \ 3,4,5,6,7,8,9,10\}\\
J_{61} &= \{\ \ 3,\quad 6,7,8,9,10\}
\end{aligned}$$

and we have $w_{\max}(\mathbf{i}(J_{61})) = s_1s_3s_2s_1s_3 = [4231] \neq w_0$, hence T is not w_0-standard.

5. Basis of $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$

Assume that $\mathbf{m}$ is regular. We shall prove that the w_0-standard monomials of shape $(\mathbf{i}, \mathbf{m})$ form a basis of the space of sections $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$. By Theorems 3.4 and 4.21, we just have to show that the restriction map

$$H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$$

is surjective. The idea is to find a sequence of varieties $\big(Y_{\mathbf{i}}^{u_t}\big)$, parametrized by $u_t \in S_n$, such that

- $Y_{\mathbf{i}}^{u_0} = Z_{\mathbf{i}}$ and $Y_{\mathbf{i}}^{u_N} = \Gamma_{\mathbf{i}}$,
- $Y_{\mathbf{i}}^{u_{t+1}}$ is a hypersurface in $Y_{\mathbf{i}}^{u_t}$,
- each restriction map $H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}})$ is surjective.

Example 5.1. Let $n = 4$ and $\mathbf{i} = 123212312$. Consider the following reduced expression

$$w_0 = s_1s_2s_1s_3s_2s_1 = s_{a_6}s_{a_5}\dots s_{a_1},$$

and set

$$\begin{cases} u_0 = e, \\ u_{t+1} = s_{a_{t+1}}u_t \quad \forall t \geq 0. \end{cases}$$

The sequence (u_t) is increasing, and $u_6 = w_0$. Thus, we obtain a sequence of opposite Schubert varieties

$$F\ell(n) = X^{u_0} \supset X^{u_1} \supset \cdots \supset X^{u_6} = \{F_{\mathrm{op\,can}}\}.$$

Let $F = (F^1 \subset F^2 \subset F^3 \subset F^4 = k^4)$ be a flag.

- We have the equivalence
$$\begin{aligned} F \in X^{s_1} &\iff F^1 \in \langle e_2, e_3, e_4 \rangle \\ &\iff p_1(F) = 0. \end{aligned}$$
So X^{u_1} is defined inside X^{u_0} by the vanishing of $p_1 = p_{\kappa_0}$.
- Assume $F \in X^{u_1}$. Then
$$\begin{aligned} F \in X^{s_2 s_1} &\iff F^1 \in \langle e_3, e_4 \rangle \\ &\iff p_2(F) = 0, \text{ since we already know that } p_1(F) = 0. \end{aligned}$$
Hence X^{u_2} is defined inside X^{u_1} by the vanishing of $p_2 = p_{\kappa_1}$.
- Similarly, X^{u_3} is defined inside X^{u_2} by the vanishing of $p_3 = p_{\kappa_2}$.
- The opposite Schubert variety X^{u_4} is defined inside X^{u_3} by the vanishing of $p_{14} = p_{\kappa_3}$.
- X^{u_5} is defined inside X^{u_4} by the vanishing of $p_{24} = p_{\kappa_4}$.
- X^{u_6} is defined inside X^{u_5} by the vanishing of $p_{134} = p_{\kappa_5}$.

We then set $Y_{\mathbf{i}}^{u_t} = \mathrm{pr}_9^{-1}(X^{u_t})$. Thus, $Y_{\mathbf{i}}^{u_0} = Z_{\mathbf{i}}$, $Y_{\mathbf{i}}^{u_6} = \Gamma_{\mathbf{i}}$. Moreover, $Y_{\mathbf{i}}^{u_{t+1}}$ is defined inside $Y_{\mathbf{i}}^{u_t}$ by the vanishing of p_{κ_t}, where we view κ_t as a tableau of shape $(123212312, \mathbf{a}'_t)$, where
$$\mathbf{a}'_1 = \mathbf{a}'_2 = \mathbf{a}'_3 = 000000010,\ \mathbf{a}'_4 = \mathbf{a}'_5 = 000000001,\ \mathbf{a}'_6 = 000000100.$$

This example leads us to work with the following varieties. Consider the last projection $\mathrm{pr}_r : Z_{\mathbf{i}} \to F\ell(n)$. Fix $u \in S_n$ and a reduced decomposition
$$w_0 u = s_{k_l} s_{k_{l-1}} \dots s_{k_1}.$$
Consider the opposite Schubert variety $X^u \subset F\ell(n)$ and set
$$Y_{\mathbf{i}}^u = \mathrm{pr}_r^{-1}(X^u) \subset Z_{\mathbf{i}}.$$
In particular, $Y_{\mathbf{i}}^e = Z_{\mathbf{i}}$ and $Y_{\mathbf{i}}^{w_0} = \Gamma_{\mathbf{i}}$.

Proposition 5.2. *The variety $Y_{\mathbf{i}}^u$ is irreducible, and if $\mathbf{i}' = i_1 \dots i_r k_1 \dots k_l$, then the projection $F\ell(n)^{r+l} \to F\ell(n)^r$ onto the r first factors restricts to a morphism*
$$\varphi : \Gamma_{\mathbf{i}'} \to Y_{\mathbf{i}}^u$$
that is birational, hence surjective.

Proof. Recall that a flag F lies in X^u if and only if it can be connected to $F_{\mathrm{op\,can}}$ by a gallery of type $k_1 \dots k_l$. Hence $Y_{\mathbf{i}}^u$ consists of all galleries
$$F_{\mathrm{can}} \overset{i_1}{\text{---}} F_1 \overset{i_2}{\text{---}} \dots \overset{i_r}{\text{---}} F_r$$
that can be extended to a gallery of the form
$$F_{\mathrm{can}} \overset{i_1}{\text{---}} F_1 \overset{i_2}{\text{---}} \dots \overset{i_r}{\text{---}} F_r \overset{k_1}{\text{---}} \dots \overset{k_l}{\text{---}} F_{\mathrm{op\,can}}.$$
Thus, φ indeed takes values in $Y_{\mathbf{i}}^u$ and is surjective. The irreducibility of $Y_{\mathbf{i}}^u$ follows.

Moreover, in the diagram
$$\begin{tikzcd} \Gamma_{\mathbf{i}'} \cong Z_{\mathbf{i}} \times_{F\ell(n)} Z_{k_l \dots k_1}(F_{\mathrm{op\,can}}) \arrow[r] \arrow[d] & Z_{k_l \dots k_1}(F_{\mathrm{op\,can}}) \arrow[d, "\mathrm{pr}_l"] \\ Z_{\mathbf{i}} \arrow[r, "\mathrm{pr}_r"'] & F\ell(n), \end{tikzcd}$$

pr_l is an isomorphism over C^u, and the morphism $\mathrm{id} \times (\mathrm{pr}_l^{-1} \circ \mathrm{pr}_r)$ from $\mathrm{pr}_r^{-1}(C^u) \subset Y_{\mathbf{i}}^u$ to $Z_{\mathbf{i}} \times Z_{k_l \dots k_1}(F_{\mathrm{op\,can}})$ is an inverse of φ over $\mathrm{pr}_r^{-1}(C^u)$, hence φ is birational. □

Corollary 5.3. *Take the notations of the previous proposition, and consider the* j*th projection* $\mathrm{pr}_j : Z_{\mathbf{i}} \to F\ell(n)$. *Then* $\mathrm{pr}_j(Y_{\mathbf{i}}^u)$ *is the Richardson variety* X_y^x *for* $y = w_{max}(i_1 \dots i_j)$ *and* $x = w_0 w_{max}(i_{j+1} \dots i_r k_1 \dots k_l)^{-1}$.

Proof. Note that $\mathrm{pr}_j(Y_{\mathbf{i}}^u) = \mathrm{pr}_j(\varphi(\Gamma_{\mathbf{i}'})) = \mathrm{pr}_j(\Gamma_{\mathbf{i}'})$ since $j \in [r]$. Proposition 1.6 leads to the result. □

Notations 5.4. As in Example 5.1, consider the reduced decomposition

$$w_0 = s_1(s_2 s_1) \dots (s_{n-1} \dots s_1) = s_{a_N} \dots s_{a_1},$$

and set $u_t = s_{a_t} \dots s_1$, $u_0 = e$.

Consider the sequence of columns κ_t defined in the following way.

- The $n-1$ first columns are $\kappa_0 = 1,\ \kappa_1 = 2, \dots,\ \kappa_{n-2} = n-1$.
- The $n-2$ next columns are $1 * n,\ 2 * n, \dots,\ n-2 * n$.
- The $n-3$ next ones are of size 3: $1 * w_0\varpi_2,\ 2 * w_0\varpi_2, \dots,\ n-3 * w_0\varpi_2$.
- We proceed in the same way for the other columns until we get $\kappa_{N-1} = 1 * w_0\varpi_{n-2}$.

We denote by b_t the size of κ_t, so that $\kappa_t = u_t\varpi_{b_t}$. We set $\kappa'_t = u_{t+1}\varpi_{b_t}$.

In the sequel, we say that a variety Y *is defined in* X *by the vanishing of* f if

$$y \in Y \iff y \in X \text{ and } f(y) = 0.$$

This does not necessarily mean that f generates the ideal of Y in the coordinate ring of X.

Lemma 5.5. *For every* $t \in [0, N-1]$, *the opposite Schubert variety* $X^{u_{t+1}} \subset F\ell(n)$ *is defined inside* X^{u_t} *by the vanishing of* p_{κ_t}.

Proof. We begin by proving the following

Claim For every t, the opposite Schubert variety $X^{u_{t+1}\varpi_{b_t}} \subset G_{b_t,n}$ is defined inside $X^{u_t\varpi_{b_t}} = X^{\kappa_t}$ by the vanishing of p_{κ_t}.

Indeed, recall that a b_t-space V belongs to the opposite Schubert variety X^{κ_t} if and only if for every $\kappa \not\geq \kappa_t$, $p_\kappa(V) = 0$, and similarly for $X^{\kappa'_t}$. Thus, we have to describe the set

$$E_t = \{\kappa \not\geq \kappa'_t \mid \kappa \geq \kappa_t\}.$$

We distinguish two cases.

- Case 1: $b_{t+1} = b_t$. Then $\kappa'_t = u_{t+1}\varpi_{b_t} = \kappa_{t+1}$. But κ_t is of the form $p * w_0\varpi_{b_t-1}$, and $\kappa_{t+1} = (p+1) * w_0\varpi_{b_t-1}$. So $\kappa_t < \kappa'_t$, hence $\kappa_t \in E_t$. Let $\kappa \in E_t$ with $\kappa \neq \kappa_t$. Then $\kappa > \kappa_t$, so $\kappa \geq \kappa_{t+1}$: a contradiction. Hence, the claim is proved in this case.
- Case 2: $b_{t+1} = b_t + 1$. Then $\kappa'_t = u_{t+1}\varpi_{b_t} = w_0\varpi_{b_t} = (n - b_t + 1) * w_0\varpi_{b_t-1}$, and $\kappa_t = (n - b_t) * w_0\varpi_{b_t-1}$. Again, $\kappa_t \in E_t$. If $\kappa \in E_t$ and $\kappa > \kappa_t$, then $\kappa = w_0\varpi_{b_t}$: a contradiction. This proves the claim.

Now, let q be the restriction to X^{u_t} of the canonical projection $F\ell(n) \to G_{b_t,n}$. We have to show that $X^{u_{t+1}} = q^{-1}(X^{\kappa'_t})$. since $u_t\varpi_{b_t} \neq \kappa'_t$, u_{t+1} is a minimal element of the poset $\{u \geq u_t,\ u\varpi_{b_t} = \kappa'_t\}$, so by Remark 4.10 $q^{-1}(X^{\kappa'_t}) = X^{u_{t+1}}$. □

Notation 5.6. For every $t \in [0, N-1]$, we set $l_t = j_{b_t}$, that is the largest integer j such that $i_j = b_t$.

Corollary 5.7. *With the notation of Lemma 5.5, the variety $Y_{\mathbf{i}}^{u_{t+1}}$ is defined inside $Y_{\mathbf{i}}^{u_t}$ by the vanishing of p_T, where $T = \emptyset * \cdots * \kappa_t * \cdots * \emptyset$ is a tableau of shape $(\mathbf{i}, 0 \dots 1 \dots 0)$, the 1 being at position l_t.*

Proof. Write $\kappa = \kappa_t$. Let γ be a gallery

$$F_{\text{can}} \overset{i_1}{—} F_1 \overset{i_2}{—} \cdots \overset{b_t}{—} F_{k_t} — \cdots — F_r$$

in $Y_{\mathbf{i}}^{u_t}$. This gallery belongs to $Y_{\mathbf{i}}^{u_{t+1}}$ if and only if $F_r \in X^{u_{t+1}}$. Since we already know that $F_r \in X^{u_t}$, we have

$$\gamma \in Y_{\mathbf{i}}^{u_{t+1}} \iff p_\kappa(F_r) = 0 \iff p_\kappa(\pi_{b_t} F_r) = 0,$$

where the first equivalence follows from Lemma 5.5 and the second from the fact that κ is of size b_t. By definition of l_t, no adjacency after F_{j_t} is an b_t-adjacency, hence $\pi_{b_t} F_{j_t} = \pi_{b_t} F_{j_t+1} = \cdots = \pi_{b_t} F_r$, and therefore,

$$p_\kappa(F_r) = 0 \iff p_\kappa(F_{j_t}) = 0 \iff p_T(\gamma) = 0,$$

where $T = \emptyset * \cdots * \kappa * \cdots * \emptyset$ with κ in position l_t. □

Notations 5.8. We fix an $\mathbf{a} = a_1 \dots a_r$ with $a_i > 0$ for every i. The associated line bundle $L_{\mathbf{i},\mathbf{a}}$ is very ample, so it induces an embedding of $Z_{\mathbf{i}}$ in some projective space $\mathbf{P}_{\mathbf{a}}$. We denote by R_t the homogeneous coordinate ring of $Y_{\mathbf{i}}^{u_t}$ viewed as a subvariety of $\mathbf{P}_{\mathbf{a}}$.

Remark 5.9. For the rest of this section, if a notion depends on an embedding, such as projective normality, or the homogeneous coordinate ring of a variety, it will be implicitly understood that we work with the line bundle $L_{\mathbf{i},\mathbf{a}}$.

The ring R_{t+1} is a quotient R_t/I_t, and we shall determine the ideal I_t. We begin by computing the equations of $Y_{\mathbf{i}}^{u_{t+1}}$ in an affine open set of $Y_{\mathbf{i}}^{u_t}$.

Definition 5.10. We shall define an affine open set Ω of $Z_{\mathbf{i}}$, isomorphic to the affine space k^r. This construction is taken from [15].

First, we define inductively a sequence of permutations (σ_j) with $\sigma_N = w_0$:

$$\begin{cases} \sigma_0 = e, \\ \sigma_{j+1} = \sigma_j * s_{i_{j+1}} \quad \forall j \geq 0. \end{cases}$$

Moreover, we set $v_{j+1} = \sigma_j^{-1} \sigma_{j+1} \in \{e, s_{i_{j+1}}\}$.

Next, consider the 1-parameter unipotent subgroup U_β associated to a root β, with its standard parametrization $\epsilon_\beta : k \to U_\beta$ (*i.e.* the matrix $\epsilon_\beta(x)$ has 1s on the diagonal, the entry corresponding to β equal to x, and 0s elsewhere). We also denote by $\alpha_1, \dots, \alpha_{n-1}$ the simple roots and by P_j the minimal parabolic subgroup associated to α_j, *i.e.* the subgroup generated by B and by $U_{-\alpha_j}$.

We set $\beta_j = v_j(-\alpha_{i_j})$ and consider the morphism

$$\begin{array}{rcl} k^r & \to & P_{\mathbf{i}} = P_{i_1} \times \dots P_{i_r} \\ (x_1, \dots, x_r) & \mapsto & (A_1, \dots, A_r) \end{array}$$

with $A_j = \epsilon_{\beta_j}(x_j) v_j$. Set $B_j = A_1 \dots A_j$.

Finally, let

$$\begin{array}{rrcl}\varphi : & k^r & \to & Z_{\mathbf{i}} \\ & (x_1, \dots, x_r) & \mapsto & (\gamma_1, \dots, \gamma_r)\end{array}$$

for $\gamma_j = B_j F_{\text{can}}$.

The image of φ is denoted by Ω: it is an open set in $Z_{\mathbf{i}}$, and $\varphi : k^r \to \Omega$ is an isomorphism.

Notation 5.11. Let $\kappa = k_1 \dots k_i$ and $\tau = t_1 \dots t_i$ be two columns of the same size. Given a matrix M, we denote by $M[\kappa, \tau]$ the determinant of the submatrix of M obtained by taking the rows $k_1, \dots, k_i$ and the columns $t_1, \dots, t_i$. Moreover, $M[\kappa, [i]]$ is simply denoted by $M[\kappa, i]$.

Example 5.12. We work on Example 5.1, where $\mathbf{i} = 123212312$, and we set $\mathbf{a} = 111111111$. The sequence (σ_j) is given by

$$\begin{array}{lll}\sigma_0 = [1234], & \sigma_1 = [2134], & \sigma_2 = [2314], \\ \sigma_3 = [2341], & \sigma_4 = [2431], & \sigma_5 = [4231], \\ \sigma_6 = [4321], & & \\ \sigma_7 = \sigma_8 = \sigma_9 = \sigma_6. & & \end{array}$$

and the sequence (v_j) is

$$\begin{array}{lll}v_1 = s_1, & v_2 = s_2, & v_3 = s_3, \\ v_4 = s_2, & v_5 = s_1, & v_6 = s_2, \\ v_7 = v_8 = v_9 = e. & & \end{array}$$

Let $T_0 = 2 * 23 * 234 * 24 * 4 * 34 * 234 * 4 * 34$. It can be shown that Ω is exactly the open set $\{\gamma \in Z_{\mathbf{i}} \mid p_{T_0}(\gamma) \neq 0\}$.

Now, direct computations show that the affine variety $Y_{\mathbf{i}}^{s_1} \cap \Omega \subset k^9$ is defined by the equation

$$Q(x_1, \dots, x_9) = x_8(x_1 x_6 + x_2) + x_1 x_5 + x_2 x_4 + x_3 = 0.$$

Since $Y_{\mathbf{i}}^{s_1} \cap \Omega$ is irreducible (as an open set of the irreducible $Y_{\mathbf{i}}^{s_1}$), this equation is also irreducible and generates the ideal of $Y_{\mathbf{i}}^{s_1} \cap \Omega$. Thus, if f is a linear combination of monomials p_T with T of shape $(\mathbf{i}, \mathbf{m})$ such that f vanishes identically on $Y_{\mathbf{i}}^{s_1}$, then $\frac{f}{T_0} \in k[x_1, \dots, x_9]$ vanishes on $Y_{\mathbf{i}}^{s_1} \cap \Omega$, hence

$$\frac{f}{T_0} \in Qk[x_1, \dots, x_9].$$

But we know that each coordinate x_j is a quotient f_j / T_0^k of degree 0 for an $f_j \in R_0 = k[Z_{\mathbf{i}}]$, and also that

$$x_8(x_1 x_6 + x_2) + x_1 x_5 + x_2 x_4 + x_3 = \frac{p_{T_1}}{p_{T_0}},$$

where $T_1 = 2 * 23 * 234 * 24 * 4 * 34 * 234 * 1 * 34$. It follows that f is a multiple of p_{T_1}, hence $f \in p_1 H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{a}'})$ where $\mathbf{a}' = 111111101$.

Hence, the ideal I_1 of $Y_{\mathbf{i}}^{u_1}$ in R_0 is $p_1 H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{a}'})$. We shall generalize this computation below.

Lemma 5.13. *For every j,*

$$U_{\beta_1} v_1 \dots U_{\beta_j} v_j = U_{\beta_1} U_{\sigma_1(\beta_2)} \dots U_{\sigma_{j-1}(\beta_j)} \sigma_j \subset B\sigma_j.$$

Proof. The equality follows from the formula

$$\sigma U_\beta = U_{\sigma(\beta)}\sigma, \quad \forall \sigma \in S_n.$$

For the inclusion, we proceed by induction over j. Since $\beta_1 = (i_1, i_1+1)$ and $v_1 = s_{i_1}$, $U_{\beta_1}v_1 \subset Bv_1 = B\sigma_1$.

Assume that the property holds for $j \geq 1$, that is $U_{\beta_1}v_1 \dots U_{\beta_j}v_j \subset B\sigma_j$.

If $\sigma_j s_{i_{j+1}} < \sigma_j$, then $\sigma_{j+1} = \sigma_j$, $v_{j+1} = e$, $\beta_{j+1} = (i_{j+1}+1, i_{j+1})$, and $\sigma_j(\beta_{j+1}) = (\sigma_j(i_{j+1}+1), \sigma_j(i_{j+1}))$.

$$\begin{aligned} \sigma_j s_{i_{j+1}} < \sigma_j &\iff \sigma_j(i_{j+1}) > \sigma_j(i_{j+1}+1) \\ &\iff U_{\sigma_j(\beta_{j+1})} \subset B. \end{aligned}$$

It follows that

$$\begin{aligned} U_{\beta_1}v_1 \dots U_{\beta_j}v_j U_{\beta_{j+1}}v_{j+1} &\subset B\sigma_j U_{\beta_{j+1}} \\ &\subset BU_{\sigma_j(\beta_{j+1})}\sigma_j \\ &\subset B\sigma_{j+1}. \end{aligned}$$

Similarly, if $\sigma_j s_{i_{j+1}} > \sigma_j$, then $\sigma_{j+1} = \sigma_j s_{i_{j+1}}$, $v_{j+1} = s_{i_j}$ and $\beta_{j+1} = \alpha_{i_{j+1}}$. Hence

$$\begin{aligned} U_{\beta_1}v_1 \dots U_{\beta_j}v_j U_{\beta_{j+1}}v_{j+1} &\subset B\sigma_j U_{\beta_{j+1}}s_{i_j} \\ &\subset BU_{\sigma_j\beta_{j+1}}\sigma_j s_{i_j} \\ &\subset B\sigma_{j+1}. \ \square \end{aligned}$$

Notation 5.14. Let κ be a column of size i. We set $O_\kappa = \{F \in F\ell(n) \mid p_\kappa(F) \neq 0\}$.

Proposition 5.15. *Let* $w \in S_n$ *and* $\kappa = w\varpi_i \in I_{i,n}$. *Then* $X_w \cap O_\kappa = \coprod_{v \in \mathcal{E}(w,\kappa)} C_v$.

Proof. Let q be the restriction to X_w of the canonical projection $F\ell(n) \to G_{i,n}$. Then we have $X_w \cap O_\kappa = q^{-1}(X_\kappa \cap q(O_\kappa))$. It is well known that $X_\kappa \cap q(O_\kappa) = C_\kappa$. But we have the equality

$$q^{-1}(C_\kappa) = \coprod_{v \in \mathcal{E}(w,\kappa)} C_v. \ \square$$

Proposition 5.16. *There exists a tableau* T_0 *of shape* $(\mathbf{i}, \mathbf{a})$ *such that*

$$\Omega = \{\gamma \in Z_{\mathbf{i}} \mid p_{T_0}(\gamma) \neq 0\}.$$

In particular, φ *induces an isomorphism* $\varphi^* : (R_0)_{(p_{T_0})} \to k[x_1, \dots, x_r]$, *where* $(R_0)_{(p_{T_0})}$ *is the subring of elements of degree 0 in the localized ring* $(R_0)_{p_{T_0}}$, i.e.

$$(R_0)_{(p_{T_0})} = \bigcup_{d \geq 0} \left\{ \frac{f}{p_{T_0}^d} \;\middle|\; f \in R_0 \text{ is homogeneous of degree } d \right\}.$$

Proof. Let $T_0 = (\sigma_1\varpi_{i_1})^{*a_1} * (\sigma_2\varpi_{i_2})^{*a_2} * \dots * (\sigma_r\varpi_{i_r})^{*a_r}$. Then

$$\{\gamma \in Z_{\mathbf{i}} \mid p_{T_0}(\gamma) \neq 0\} = \prod_{j=1}^{r} \mathrm{pr}_j^{-1}\left(O_{\sigma_j\varpi_{i_j}}\right).$$

We know that

$$\mathrm{pr}_j(\Omega) = U_{\beta_1}v_1 \dots U_{\beta_j}v_j F_{\mathrm{can}}.$$

Thus, by Lemma 5.13, $\mathrm{pr}_j(\Omega) \subset B\sigma_j F_{\mathrm{can}} = C_{\sigma_j}$. But if $F \in C_{\sigma_j}$, then its i_j-th constituent F^{i_j} belongs to $C_{\sigma_j\varpi_{i_j}}$, so

$$p_{\sigma_j\varpi_{i_j}}(F) = p_{\sigma_j\varpi_{i_j}}(F^{i_j}) \neq 0.$$

This proves the inclusion

$$\Omega \subset \prod_{j=1}^{r} \mathrm{pr}_j^{-1}\left(O_{\sigma_j \varpi_{i_j}}\right).$$

For the opposite inclusion, we proceed by induction over r.

If $r = 1$, then $\Omega = \{F_{\mathrm{can}}\} \times U_{\beta_1} v_1 F_{\mathrm{can}}$, with $v_1 = s_{i_1}$ and $\beta_1 = \alpha_{i_1}$. Hence $\Omega = \{F_{\mathrm{can}}\} \times C_{s_{i_1}}$. By Lemma 5.15, $X_{s_{i_1}} \cap O_{s_{i_1}\varpi_{i_1}} = C_{s_{i_1}}$, so

$$\mathrm{pr}_1^{-1}(O_{\sigma_1 \varpi_{i_1}}) = \{F_{\mathrm{can}} \overset{i_1}{\text{---}} F^1 \mid F^1 \in C_{s_{i_1}}\}.$$

Thus $\Omega = \mathrm{pr}_1^{-1}(O_{\sigma_1\varpi_{i_1}})$.

Let $r > 1$ and assume that the property holds for $r-1$. Let $\gamma \in \prod_{j=1}^{r} \mathrm{pr}_j^{-1}\left(O_{\sigma_j \varpi_{i_j}}\right)$. By induction, there exist $A_1, \ldots, A_{r-1}$ such that $A_j \in U_{\beta_j} v_j$ and $\gamma_j = A_1 \ldots A_j F_{\mathrm{can}}$ for $j \leq r-1$. Since $\gamma_{r-1} \overset{i_r}{\text{---}} \gamma_r$, there exists $p \in P_{i_r}$ such that $A_1 \ldots A_{r-1} p F_{\mathrm{can}} = \gamma_r$. Now, $P_{i_r} F_{\mathrm{can}}$ is a T-stable curve and we have

$$P_{i_r} F_{\mathrm{can}} = U_{-\alpha_{i_r}} F_{\mathrm{can}} \cup \{s_{i_r} F_{\mathrm{can}}\} = U_{\alpha_{i_r}} s_{i_r} F_{\mathrm{can}} \cup \{F_{\mathrm{can}}\}.$$

If $v_r = e$, then $\sigma_r = \sigma_{r-1}$ and $\beta_r = -\alpha_{i_r}$. By Lemma 5.13, $\gamma_r \in B\sigma_{r-1} p F_{\mathrm{can}}$. If $pF_{\mathrm{can}} = s_{i_r} F_{\mathrm{can}}$, then $\gamma_r \in C_{\sigma_r s_r}$. But by Lemma 5.15, γ_r belongs to a Schubert cell C_v with $v\varpi_{i_r} = \sigma_r \varpi_{i_r}$. Since $\sigma_r \varpi_{i_r} \neq \sigma_r s_r \varpi_{i_r}$, we have a contradiction. Hence $pF_{\mathrm{can}} \in U_{-\alpha_{i_r}} F_{\mathrm{can}}$. So we may choose p in U_{β_r}, which proves that $\gamma \in \Omega$.

If $v_r = s_{i_r}$, we prove similarly that $pF_{\mathrm{can}} \neq F_{\mathrm{can}}$, so we may choose p in $U_{\alpha_{i_r}} s_{i_r} = U_{\beta_r} v_r$, thus $\gamma \in \Omega$. □

Remark 5.17. Consider an arbitrary tableau T of shape $(\mathbf{i}, \mathbf{a})$. Then we may compute $\varphi^*\left(\frac{p_T}{p_{T_0}}\right)$ in the following way. Write

$$T = \kappa_{11} * \cdots * \kappa_{1a_1} * \cdots * \kappa_{r1} * \cdots * \kappa_{ra_r},$$

then

$$\varphi^*\left(\frac{p_T}{p_{T_0}}\right)(x_1, \ldots, x_r) = B_1[\kappa_{11}, i_1] \ldots B_1[\kappa_{1a_1}, i_1] \ldots B_r[\kappa_{r1}, i_r] \ldots B_r[\kappa_{ra_r}, i_r].$$

Proposition 5.18. *Denote by $Q_{ij} \in k[x_1, \ldots, x_r]$ the entries of B_r:*

$$B_r = \begin{pmatrix} Q_{1,1} & Q_{2,1} & \cdots & \cdots & Q_{n-1,1} & 1 \\ \vdots & \vdots & & & 1 & 0 \\ \vdots & \vdots & & \iddots & 0 & 0 \\ \vdots & \vdots & \iddots & \iddots & \vdots & \vdots \\ Q_{n-1,1} & 1 & \iddots & & \vdots & \vdots \\ 1 & 0 & \cdots & \cdots & 0 & 0 \end{pmatrix}.$$

The polynomials Q_{ij} all have distinct linear parts.

Proof. It suffices to prove that $B_r w_0 \in B$, but this follows from Lemma 5.13:

$$B_r \in U_{\beta_1} v_1 \ldots U_{\beta_r} v_r F_{\mathrm{can}} \subset B\sigma_r = Bw_0.$$

We may obtain the linear part of the Q_{ij} by derivating B_r. From the expression $B_r = \epsilon_{\beta_1}(x_1)v_1 \dots \epsilon_{\beta_r}(x_r)v_r$, we see that

$$\frac{\partial B_r}{\partial x_j}(0) = E_{\sigma_{j-1}(\beta_j)} w_0.$$

This already proves that the linear parts of the Q_{ij} are distinct. We shall now prove that every elementary matrix E_{kl} occurs as a derivative of B_r, that is, each pair $(i, i+1)$ equals some $\sigma_{j-1}(\beta_j)$, or equivalently, $U_{\beta_1} U_{\sigma_1 \beta_2} \dots U_{\sigma_{r-1}\beta_r} = U$ (where U is the unipotent part of B). Since $\mathrm{pr}_r(\Omega) = C_{w_0}$, we have

$$U_{\beta_1} U_{\sigma_1(\beta_2)} \dots U_{\sigma_{r-1}(\beta_r)} F_{\mathrm{can}} = U_{\beta_1} v_1 \dots U_{\beta_r} v_r F_{\mathrm{can}} = U w_0 F_{\mathrm{can}}.$$

Since the stabilizer of $w_0 F_{\mathrm{can}}$ in U is trivial, we conclude that $U_{\beta_1} U_{\sigma_1 \beta_2} \dots U_{\sigma_{r-1}\beta_r} = U$. $\square$

Proposition 5.19. *The ideal of* $Y_{\mathbf{i}}^{u_{t+1}} \cap \Omega$ *in the coordinate ring of* $Y_{\mathbf{i}}^{u_t} \cap \Omega$ *is generated by* Q_{κ_{t1}, b_t}*, where* κ_{t1} *is the first entry of the column* κ_t*. Moreover,*

$$Q_{\kappa_{t1}, b_t} = \varphi^* \left(\frac{p_{T_t}}{p_{T_0}} \right)$$

where T_t *is the tableau obtained from* T_0 *by replacing its last column of size* b_t *by* κ_t*.*

Moreover, the varieties $Y_{\mathbf{i}}^{u_t} \cap \Omega$ *are isomorphic to affine spaces.*

Proof. We alredady know that $Y_{\mathbf{i}}^{u_{t+1}}$ is defined inside $Y_{\mathbf{i}}^{u_t}$ by the vanishing of p_{κ_t}: given a gallery $\gamma = (F_{\mathrm{can}} \overset{i_1}{\text{—}} F_1 \overset{i_2}{\text{—}} \dots \overset{i_r}{\text{—}} F_r)$ in $Y_{\mathbf{i}}^{u_t}$, we know by Corollary 5.7 that $\gamma \in Y_{\mathbf{i}}^{u_{t+1}}$ if and only if $p_{\kappa_t}(F_{l_t}) = 0$.

In Ω, this corresponds to the vanishing of $B_{l_t}[\sigma_{l_t} \varpi_{b_t}, b_t]$. Now, as in the proof of Lemma 5.13,

$$B_r = B_{l_t} b v_{l_{t+1}} \dots v_r$$

for some $b \in U$. The jth column of $B_{l_t} b$ is then a linear combination of the columns $1, \dots, j$ of B_{l_t}. So

$$(B_{l_t} b)[\sigma_{l_t} \varpi_{b_t}, b_t] = B_{l_t}[\sigma_{l_t} \varpi_{b_t}, b_t].$$

Moreover, by definition of l_t, the permutation $v_{l_{t+1}} \dots v_r$ stabilizes the fundamental weight column ϖ_{b_t}, so $B_{l_t} b$ and B_r have the same first b_t columns up to a permutation, hence

$$B_{l_t}[\sigma_{l_t} \varpi_{b_t}, b_t] = \pm B_r[\sigma_{l_t} \varpi_{b_t}, b_t].$$

A straightforward computation shows that this determinant is $\pm Q_{\kappa_{t1}, b_t}$.

To prove that $\varphi^* \left(\frac{p_{T_t}}{p_{T_0}} \right) = \pm Q_{\kappa_{t1}, b_t}$, note that

$$\varphi^* \left(\frac{p_{T_t}}{p_{T_0}} \right) = B_1[\sigma_1 \varpi_{i_1}, i_1] \dots B_{j_t}[\sigma_{j_t} \varpi_{b_t}, b_t] \dots B_r[\sigma_r \varpi_{i_r}, i_r].$$

Now, by Lemma 5.13, $B_j = b_j \sigma_j$ for some $b_j \in B$. So, for $j \neq l_t$,

$$B_j[\sigma_j \varpi_{i_j}, i_j] = \pm b_j[\sigma_j \varpi_{i_j}, \sigma_j \varpi_{i_j}] = \pm 1.$$

Hence $Y_{\mathbf{i}}^{u_{t+1}} \cap \Omega$ is defined by the equation $Q_{\kappa_{t1}, b_t} = 0$. But this polynomial is of the form $x_{p_t} - Q'$ for some variable $x_{p_t} \in \{x_1, \dots, x_r\}$, so we may substitute x_{p_t} by Q' in the coordinate ring of $Y_{\mathbf{i}}^{u_t} \cap \Omega$ to obtain the coordinate ring of $Y_{\mathbf{i}}^{u_{t+1}} \cap \Omega$. Thus, by induction over t, we see that the coordinate ring of $Y_{\mathbf{i}}^{u_{t+1}} \cap \Omega$ is isomorphic to $k[x_i \mid i \neq p_0, \dots, p_t]$. In particular, this ring is a Unique Factorization Domain.

Therefore, the irreducible polynomial Q_{κ_{t1},b_t} generates the ideal of $Y_{\mathbf{i}}^{u_{t+1}}$ in the coordinate ring of $Y_{\mathbf{i}}^{u_t}$. □

Notations 5.20. We set $\mathbf{a}'_t = 0 \dots -1 \dots 0$, the -1 again being at position l_t. Let S_t be the R_t-graded module associated to the coherent sheaf $L_{\mathbf{i},\mathbf{a}'_t}$, that is,

$$S_t = \bigoplus_{d=0}^{+\infty} H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},d\mathbf{a}+\mathbf{a}'_t}).$$

Assumption 5.21. We assume that $\mathbf{a} + \sum_{t=0}^{N-1} \mathbf{a}'_t$ is regular.

Corollary 5.22. *Denote by $\mathcal{O}_{Y_{\mathbf{i}}^{u_t}}$ the structural sheaf of $Y_{\mathbf{i}}^{u_t}$ and assume that $Y_{\mathbf{i}}^{u_t}$ is projectively normal. Then the sequence of R_t-modules*

$$(*) \qquad 0 \to S_t \to R_t \to R_{t+1} \to 0$$

is exact, where the first map is the multiplication by p_{κ_t} and the second is the natural projection.

The exact sequence $(*)$ induces an exact sequence of sheaves of $\mathcal{O}_{Y_{\mathbf{i}}^{u_t}}$-modules

$$0 \to L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t} \to L_{\mathbf{i},\mathbf{m}} \to (L_{\mathbf{i},\mathbf{m}})_{|Y_{\mathbf{i}}^{u_{t+1}}} \to 0$$

and a long exact sequence in cohomology

$$(**) \qquad \begin{tikzcd}
0 \arrow[r] & H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t}) \arrow[r] & H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) \arrow[r] & H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \arrow[dll, rounded corners] \\
 & H^1(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t}) \arrow[r] & \cdots & \\
 & & \cdots \arrow[r] & H^{i-1}(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \arrow[dll, rounded corners] \\
 & H^i(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t}) \arrow[r] & H^i(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) \arrow[r] & H^i(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \arrow[dll, rounded corners] \\
 & H^{i+1}(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t}) \arrow[r] & \cdots &
\end{tikzcd}$$

Proof. Since $Y_{\mathbf{i}}^{u_t}$ is projectively normal, we know that

$$R_t = \bigoplus_{d=0}^{+\infty} H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},d\mathbf{a}}),$$

hence the sequence

$$\begin{tikzcd}
0 \arrow[r] & S_t \arrow[r, "\mu"] & R_t \arrow[r, "q"] & R_{t+1} \arrow[r] & 0
\end{tikzcd}$$

is well defined. Moreover, we already know by Corollary 5.7 that $q \circ \mu = 0$.

Let f be an homogeneous element of degree d in R_t, and suppose that $q(f) = 0$, that is, f vanishes identically on $Y_{\mathbf{i}}^{u_{t+1}}$. Then $\frac{f}{p_{T_0}^d}$ vanishes identically on $Y_{\mathbf{i}}^{u_{t+1}} \cap \Omega$, hence $\varphi^*\left(\frac{f}{p_{T_0}^d}\right) \in k[x_1,\ldots,x_r]$ is a multiple of $Q_{\kappa_{t1},b_t} = \varphi^*\left(\frac{p_{T_t}}{p_{T_0}}\right)$. It follows that f is a multiple p_{T_t}, hence $f \in p_{\kappa_t}S_t = \mu(S_t)$.

If we consider the coherent sheaves associated to these R_t-modules and tensor them by $L_{\mathbf{i},\mathbf{m}}$, then we get the exact sequence of sheaves of $\mathcal{O}_{Y_{\mathbf{i}}^{u_t}}$-modules

$$0 \to L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t} \to L_{\mathbf{i},\mathbf{m}} \to (L_{\mathbf{i},\mathbf{m}})_{|Y_{\mathbf{i}}^{u_{t+1}}} \to 0,$$

which gives the long exact sequence $(**)$. □

Theorem 5.23.

$(1)_t$ *For every t, the variety $Y_{\mathbf{i}}^{u_t}$ is projectively normal.*

$(2)_t$ *For every $i > 0$, and every $\mathbf{m}$ such that $\mathbf{m}_t = \mathbf{m} + \mathbf{a}'_0 + \cdots + \mathbf{a}'_{t-1}$ is regular, $H^i(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) = 0$. In particular, $H^i(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},d\mathbf{a}}) = 0$ for every $d \in \mathbf{Z}_{\geq 0}$.*

$(3)_t$ *If $t \geq 0$ and $\mathbf{m}_{t+1}$ is regular, then the restriction map $H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}})$ is surjective.*

Proof. We proceed by induction over t.

For $t = 0$, $Y_{\mathbf{i}}^{u_0} = Z_{\mathbf{i}}$. By Theorem 2.6, $Z_{\mathbf{i}}$ is projectively normal, and $H^i(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) = 0$ for $i > 0$. Assume now that $\mathbf{m}_1 = \mathbf{m} + \mathbf{a}'_0$ is regular. Since $Z_{\mathbf{i}}$ is projectively normal, by Corollary 5.22, we have the exact sequence

$$0 \to H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}_1}) \to H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_1}, L_{\mathbf{i},\mathbf{m}}) \to H^1(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}_1}).$$

Since $H^1(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}_1}) = 0$, the restriction map $H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_1}, L_{\mathbf{i},\mathbf{m}})$ is surjective.

Assume that the theorem is true for a $t \geq 0$.

We shall prove that $(1)_{t+1}$ is true. By induction, $Y_{\mathbf{i}}^{u_t}$ is projectively normal, so the sequence

$$0 \to H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t}) \to H^0(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \to H^1(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t})$$

is exact. Moreover, for $d \in \mathbf{Z}_{\geq 0}$, by $(2)_t$ we have $H^1(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},d\mathbf{a}}) = 0$, hence an exact sequence

$$0 \to S_t \to R_t \to \bigoplus_{d\geq 0} H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},d\mathbf{a}}) \to 0.$$

Since the sequence

$$0 \to S_t \to R_t \to R_{t+1} \to 0$$

is also exact, we have $R_{t+1} = \bigoplus_{d\geq 0} H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},d\mathbf{a}})$, that is, $Y_{\mathbf{i}}^{u_{t+1}}$ is projectively normal.

We now prove that $(2)_{t+1}$ is true. Let $\mathbf{m}$ be such that $\mathbf{m}_{t+1}$ is regular. Note that $\mathbf{m}_t$ is also regular. Since $Y_{\mathbf{i}}^{u_t}$ is projectively normal, we have the exact sequence

$$H^i(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) \to H^i(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \to H^{i+1}(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t}).$$

Now, by $(2)_t$,

$$\begin{aligned} \mathbf{m}_t \text{ regular} &\implies H^i(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}}) = 0, \\ \mathbf{m}_{t+1} \text{ regular} &\implies H^{i+1}(Y_{\mathbf{i}}^{u_t}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_t}) = 0. \end{aligned}$$

Thus, $H^i(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) = 0$.

Finally, we prove $(3)_{t+1}$. Since $Y_{\mathbf{i}}^{u_{t+1}}$ is projectively normal, we have the exact sequence

$$0 \to H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_{t+1}}) \to H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_{t+2}}, L_{\mathbf{i},\mathbf{m}}) \to H^1(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_{t+1}}).$$

Since $\mathbf{m}_{t+2}$ is regular, we have by $(2)_{t+1}$ that $H^1(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}+\mathbf{a}'_{t+1}}) = 0$, hence the restriction map $H^0(Y_{\mathbf{i}}^{u_{t+1}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(Y_{\mathbf{i}}^{u_{t+2}}, L_{\mathbf{i},\mathbf{m}})$ is surjective. □

Corollary 5.24. *If* $\mathbf{m}_N = \mathbf{m} + \sum_{t=0}^{N-1} \mathbf{a}'_t$ *is regular, then a basis of* $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$ *is given by the* w_0*-standard monomials of shape* $(\mathbf{i}, \mathbf{m})$.

Proof. Since the restriction $H^0(Z_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}}) \to H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$ is surjective, the standard monomials p_T that do not vanish identically on $\Gamma_{\mathbf{i}}$ form a generating set. By Theorem 4.21, these monomials are exactly the w_0-standard monomials. By Theorem 3.4, these monomials are linearly independent. □

Proposition 5.25. *Let* p_T *be a standard monomial of shape* $(\mathbf{i}, \mathbf{m})$*, with* $\mathbf{m}$ *arbitrary. Then* p_T *decomposes as a linear combination of* w_0*-standard monomials on* $\Gamma_{\mathbf{i}}$.

Proof. With the notation of Theorem 4.21, the result is true if $\mathbf{m}_N$ is regular. If this is not the case, then we set $\mathbf{b} = b_1 \dots b_r$ with

$$b_j = \begin{cases} N & \text{if } j = l_t \text{ for some } t, \\ 0 & \text{otherwise,} \end{cases}$$

and $\mathbf{m}' = \mathbf{m} + \mathbf{b}$. Now, $\mathbf{m}'$ satisfies the assumption of Theorem 4.21. We multiply p_T by $p_{\kappa_1}^N \dots p_{\kappa_{n-1}}^N$ to obtain a new monomial p'_T, where $\kappa_j = w_0\varpi_j$. So p'_T is of shape $(\mathbf{i}, \mathbf{m}')$ and does not vanish identically on $\Gamma_{\mathbf{i}}$. Now, p'_T may be non standard, so we decompose it as a linear combination of w_0-standard monomials of shape $(\mathbf{i}, \mathbf{m}')$ on $\Gamma_{\mathbf{i}}$, thanks to Corollary 5.24:

$$p_T(p_{\kappa_1} \dots p_{\kappa_{n-1}})^N = p_{T'} = \sum_{T''} a_{T''} p_{T''}.$$

Since a w_0-standard monomial does not vanish identically on $\Gamma_{\mathbf{i}}$, the columns κ that are in position j_k in a tableau T'' are maximal, *i.e.* equal to $w_0\varpi_k$. Hence we may factor this linear combination by $(p_{\kappa_1} \dots p_{\kappa_{n-1}})^N$, so that p_T is a linear combination of w_0-standard monomials. □

Corollary 5.26. *A basis of* $H^0(\Gamma_{\mathbf{i}}, L_{\mathbf{i},\mathbf{m}})$ *is given by the* w_0*-standard monomials of shape* $(\mathbf{i}, \mathbf{m})$. □

Remark 5.27. In the regular case ($m_i > 0$ for every i), the basis given by standard monomials is compatible with $\Gamma_{\mathbf{i}}$: this is no longer the case if $\mathbf{m}$ is not regular, see Remark 4.22.

REFERENCES

[1] M. Balan, *Multiplicity on a Richardson variety in a cominuscule G/P*, to appear in Trans. Amer. Math. Soc.; Preprint `http://arxiv.org/abs/1009.2873`.

[2] S. Billey, I. Coskun, *Singularities of generalized Richardson varieties*, Preprint `http://arxiv.org/abs/1008.2785`.

[3] A. Björner, F. Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, 231, Springer, New-York, 2005.

[4] R. Bott, H. Samelson, *Applications of the theory of Morse to symmetric spaces*, Amer. J. Math. **80** (1958), 964–1029.

[5] M. Brion, *Positivity in the Grothendieck group of complex flag varieties*, J. Algebra **258** (2002), no. 1, 137–159.

[6] M. Brion, *Lectures on the geometry of flag varieties*, Topics in cohomological studies of algebraic varieties, 33–85, Trends Math., Birkhäuser, Basel, 2005.

[7] M. Brion, V. Lakshmibai, *A geometric approach to standard monomial theory*, Represent. Theory **7** (2003), 651–680 (electronic).

[8] C. Contou-Carrère, *Géométrie des groupes semi-simples, résolutions équivariantes et lieu singulier de leurs variétés de Schubert*, Thèse d'État, Université de Montpellier II, 1983.

[9] M. Demazure, *Désingularisation des variétés de Schubert généralisées*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 53–88.

[10] V. Deodhar, *On some geometric aspects of Bruhat orderings. I. A finer decomposition of Bruhat cells*, Invent. Math. **79** (1985), no. 3, 499–511.

[11] P. Foth, S. Kim, *Standard Monomial Theory of RR varieties*, Preprint `http://arxiv.org/abs/1009.1645`.

[12] S. Gaussent, *The fibre of the Bott-Samelson resolution*, Indag. Math. (N.S.) **12** (2001), no. 4, 453–468.

[13] K. Goodearl, M. Yakimov, *Poisson structures on affine spaces and flag varieties. II*, Trans. Amer. Math. Soc. **361** (2009), no. 11, 5753–5780.

[14] H. Hansen, *On cycles in flag manifolds*, Math. Scand. **33** (1973), 269–274.

[15] M. Härterich, *The T-equivariant Cohomology of Bott-Samelson varieties*, Preprint `http://arxiv.org/abs/math/0412337`.

[16] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52, Springer-Verlag, New York-Heidelberg, 1977.

[17] X. He, T. Lam, *Projected Richardson varieties and affine Schubert varieties*, Preprint `http://arxiv.org/abs/1106.2586`.

[18] X. He, J. Lu, *On intersections of certain partitions of a group compactification*, Int. Math. Res. Not. 2011, no. 11, 2534–2564.

[19] W. V. D. Hodge, *Some enumerative results in the theory of forms*, Proc. Cambridge Philos. Soc. **39** (1943). 22–30.

[20] W. Hodge, D. Pedoe, *Methods of algebraic geometry. Vol. II*, Cambridge University Press, Cambridge, 1952.

[21] A. Knutson, T. Lam, D. Speyer, *Positroid varieties I: Juggling and Geometry*, Preprint `http://arxiv.org/abs/0903.3694`.

[22] A. Knutson, T. Lam, D. Speyer, *Projections of Richardson Varieties*, Preprint `http://arxiv.org/abs/1008.3939`.

[23] V. Kreiman, *Local properties of Richardson varieties in the Grassmannian via a bounded Robinson-Schensted-Knuth correspondance*, J. Algebraic Combin. **27** (2008), no. 3, 351–382.

[24] V. Kreiman, V. Lakshmibai, *Richardson Varieties in the Grassmannian*, in "Contributions to automorphic forms, geometry, and number theory", 573–597, Johns Hopkins Univ. Press, Baltimore, MD, 2004.

[25] V. Lakshmibai, P. Littelmann, *Richardson varieties and equivariant K-theory*, J. Algebra **260** (2003), no. 1, 230–260.

[26] V. Lakshmibai, P. Littelmann, P. Magyar, *Standard Monomial Theory and Applications. Notes by Rupert W. T. Yu*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., **514**, Representation theories and algebraic geometry (Montreal, PQ, 1997), 319–364, Kluwer Acad. Publ., Dordrecht, 1998.

[27] V. Lakshmibai, P. Littelmann, P. Magyar, *Standard monomial theory for Bott-Samelson varieties*, Compositio Math. **130** (2002), no. 3, 293–318.

[28] V. Lakshmibai, P. Magyar, *Standard monomial theory for Bott-Samelson varieties of* $\mathrm{GL}(n)$, Publ. Res. Inst. Math. Sci. **34** (1998), no. 3, 229–248.

[29] V. Lakshmibai, C. S. Seshadri, *Geometry of G/P. V*, J. Algebra **100** (1986), no. 2, 462–557.

[30] V. Lakshmibai, C. S. Seshadri, *Standard monomial theory*, Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989), 279–322, Manoj Prakashan, Madras, 1991.

[31] N. Lauritzen, J. Thomsen, *Line bundles on Bott-Samelson varieties*, J. Algebraic Geom. **13** (2004), no. 3, 461–473.

[32] P. Littelmann, *A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras*, Invent. Math. **116** (1994), no. 1-3, 329–346.

[33] P. Littelmann, *Paths and root operators in representation theory*, Ann. of Math. (2) **142** (1995), no. 3, 499–525.

[34] P. Littelmann, *Contracting modules and standard monomial theory for symmetrizable Kac-Moody algebras*, J. Amer. Math. Soc. **11** (1998), no. 3, 551–567.

[35] P. Magyar, *Schubert polynomials and Bott-Samelson varieties*, Comment. Math. Helv. **73** (1998), no. 4, 603–636.

[36] R. Richardson, *Intersections of double cosets in algebraic groups*, Indag. Math. (N.S.) **3** (1992), no. 1, 69–77.

[37] R. Richardson, T. Springer, *Combinatorics and geometry of K-orbits on the flag manifold*, Linear algebraic groups and their representations (Los Angeles, CA, 1992), 109–142, Contemp. Math., 153, Amer. Math. Soc., Providence, RI, 1993.

[38] R. Stanley, *Some combinatorial aspects of the Schubert calculus*, Combinatoire et représentation du groupe symétrique (Actes Table Ronde CNRS, Univ. Louis-Pasteur Strasbourg, Strasbourg, 1976), pp. 217–251. Lecture Notes in Math., Vol. **579**, Springer, Berlin, 1977.

[39] B. Taylor, *A straightening algorithm for row-convex tableaux*, J. Algebra **236** (2001), no. 1, 155–191.

[40] S. Upadhyay, *Initial ideals of tangent cones to Richardson varieties in the Orthogonal Grassmannian via a Orthogonal-Bounded-RSK-Correspondence*, Preprint `http://arxiv.org/abs/0909.1424`.

Université de Valenciennes, Laboratoire de Mathématiques, Le Mont Houy – ISTV2, F-59313 Valenciennes Cedex 9, France

E-mail address: `michael.balan@univ-valenciennes.fr`